INDEX FOR VOLUME 43

AUTHOR AND SUBJECT INDEX. New species and varieties in bold face type. Junior authorship indicated by pages in parentheses.

Abaca sheath rot, (*Marasmius stenophyllus*) susceptibility and control, 479
ABBOTT, E. V., 289
Acer macrophyllum, apple flyspeck host, 586
Acidity, see Hydrogen-ion concentration
Actidione, see under Antibiotics
Actinomycete, gummosis of grapefruit, associated with, 101
ADAM, ANGELOS, 465
ADAMS, D. F., (587)
ADAMS, ROBERT E., 465
Adenine, effect on rooting of *Prunus mahaleb*, and effect of pretreatment on effect of, 587
ADP, 224
Aegilops triuncialis, 310
African violet, root-knot nematode, 406
Agallia constricta, see under Insects
Agalliopsis novella, see under Insects
Agaricaceae, wood rotting by, 21
Agrilus bilineatus, see under Insects
Agrobacterium tumefaciens,
  colonial variants, morphology, 509
    pathogenicity and response to biochemical tests, 510
    stability in galls and culture, 510
Agropyron trachycaulum, 309
AKELEY, ROBERT V., (245)
Albugo candida, overwintering, 485
Aldehyde, effect on virulence of Pythium root-rot pathogen, 289
Alfalfa black stem, 378
    control, fungicides, 379
    effect on seed crop in field and greenhouse tests, 378
    effect on seed germination, 381
    measurement of seed loss from, 380
  crown rot, hydrogen cyanide production by pathogen, effect of temperature and media compared, 581
  leaf spot, resistance, classification of plants for, 653
    microscopic evidences of resistance, 652
    rapid method of measurement of resistance, 651
    Pseudoplea, Stemphylium botryosum as perfect stage, 584
    Stemphylium, 477
  mosaic virus, complex, 271
    strain, age of plant and method of inoculation, 473
    strains, 275
      differentiation, 479
    tomato varieties, 479
  nurse plant for leafhopper vectors of wound-tumor virus, 387
  Pierce's disease, grape degeneration on, 611
    yellow-headed sharpshooter vector, 612
  Pythium spp. pathogenicity on seedlings, 474
  root-rot, Phytophthora, 586
    tests for control, 589
  winter crown rot, HCN production by pathogen, 478
    host parasite relation, 478
  wound-tumor virus, insect vectors, 9
  yellow mosaic virus (*Marmor medicaginis* H. var. **flavovarians** n. var.) on, 38
ALLEN, PAUL J., 221
ALLEN, ROSS M., 585
Allium cepa, see Onion
Almond, *Sclerotinia fructicola*, and *S. laxa*, pathogenicity compared, 589
Alternaria porri, onion, effect on in storage, 411
    onion purple and brown blotch, 409
  solani, inhibition by phosphate buffer, 478
    nutrition of host, effect on susceptibility to, 476
    toximycin as protectant spray, 25
  sp., in bean seeds, 483
Alternaria—*continued*
    spp., on peanuts in market, 474
      on soybean, 292
  tenuis and *tenuissima*, 409, 411
Althaea rosea, see hollyhock
Amaryllis, spiral nematode, 297
American Phytopathological Society:
  abstracts for the 45th annual meeting (Madison), 465
  analysis of membership, D1
  charter members, D4
  classes of membership, D1
  constitution, D55
  directory of members, D10
  divisions, D2
  fields of interest reported by members, D3
  geographical index of members, D46
  growth, D1
  in memorium, D6
  introduction to directory, D1
  organization, 1953, D9
  past councilors, D7
  past editors-in-chief, D7
  past officers, D7
  professions and occupations of members, D2
  report of 44th annual meeting, 52
  reports and abstracts:
    Northeastern Division, 108
    Pacific Division, 585
    Potomac Division, 404
    Southern Division, 288
  source of employment of members, D2
  sustaining associates, D10
Amino acids, see also under Nutrition of microorganisms
  carnation bacterial wilt studies, 587
  effect on melanin production and growth in *Streptomyces scabies*, 357
  production by *Ustilago zeae* in culture, 470
Ammate, wheat stem rust, 661
ANDERSON, E. J., 465
ANDERSON, H. W., (465)
Andropogon barbinodis, 310
Annulus, see also Viruses
  rhubarb ringspot-like virus, 590
Antagonisms, *Fusarium vasinfectum*, soil microorganisms, 482
  Penicillium soil isolate to fungi, 479
  soil microflora, *Sclerotium rolfsii*, 366
Anthoxanthum odoratum, 310
Anthracnose, oak, 103
  watermelon, breeding for resistance to, 254
Antibiotic production, by *Aspergillus clavatus* and *Penicillium patulum*, 108
Antibiotics, actidione, 486
    bordeaux compared, cherry leaf spot, 468
    cherry leaf spot, 109, 468
    Curvularia "fading-out" on turf, 109
    lime, effect upon effectiveness, 109
    methoxychlor, compatible with, 109
    orthocide mixture, wheat covered smut, 461
    pH effect on activity, wheat rusts, 498
    Rhizoctonia, 475
    wheat leaf and stem rust, control and phytotoxicity, 493
  antimycin A,
    A-102, A-35 composition compared, 478
      oat seedling blight, 478
      produced by new isolate of *Streptomyces* sp., 478
      tomato early blight protectant, 478
    effect on respiration, 227
  aureomycin, oxidative phosphorylations, action on, 225
    respiration, effect on, 221
    Verticillium, 475
  bacitracin, *Rhizoctonia*, 475

Antibiotics—*continued*
chloramphenicol, effect of concentration in medium on synthesis by *Streptomyces venezuelae*, 478
chloromycetin, *Rhizoctonia*, 475
gramicidin, oxidative phosphorylations, action on, 225
respiration, effect on, 221
Staphylococcus aureus, reaction to, 225
helixin B, action across air space on *Glomerella cingulata* spores, 393
barley seedling blight, field and greenhouse tests, 392
oat Helminthosporium blight in field and greenhouse tests, 391
seed treatment, small grain, 391
inhibition of, 233
magnamycin, *Rhizoctonia* and *Verticillium*, 475
naturally occurring, 226
patulin and vivotoxins, 233
penicillin, *Rhizoctonia*, 475
polymixin, *Rhizoctonia* and *Verticillium*, 475
potato late blight effect on, 463
respiration, effect on, in microorganisms, 225
in plant tissue, 221
rimocidin, potato late blight, 464
Rhizoctonia, 475
streptomycin, bean halo blight, 407, 480
effect on Bartlett pear seedlings, 405
effect on Elberta peach, 405
effect on respiration, 227
in isolation of *Fusarium* spp. from plant tissues, 700
movement in bean, 480
Rhizoctonia and *Verticillium*, 475
terramycin, effect of concentration in medium upon synthesis by *Streptomyces venezuelae*, 478
effect on Elberta peach, 405
peach leaf drop, 405
potato late blight, 464
Verticillium, 475
thiolutin, potato late blight, 464
Rhizoctonia and *Verticillium*, 475
toximycin, antifungal, from *Bacillus subtilis*, 23
usnic acid, oxidative phosphorylations, action on, 225
respiration, effect on, 221
Antimycin, see under Antibiotics
Antirrhinum majus, see Snapdragon
Aphanomyces cochlioides, sugar beet seedlings, black rot, 300
Aphids, see under Insects
Apium graveolens, see also Celery
bayberry-yellows virus on, 18
western mosaic aphid transmission, 29
APPLE, J. L., 289
Apple, black rot, effect of continual use of fungicides, 268
Botryosphaeria rot pathogens, susceptible apple varieties, 407
disease control, mist and hydraulic spraying compared, 465
Dothiorella spp. associated with cankered branches, 407
fluoride concentrations, foliar, factors affecting, 587
flyspeck, ecology and host range, 586
perfect and imperfect stage of pathogen, 470
scab, control by fungicide mixtures, 109
seedlings in greenhouse, testing of fungicides on, 109
Apricot, *Sclerotinia fructicola* and *S. laxa*, pathogenicity compared, 589
Arachis hypogaea, see Peanut
ARAGAKI, M., (468), 643
Arbutus menziesii, apple flyspeck host, 586
Arctium minus, *Verticillium albo-atrum* on, 588
Aristida oligantha, 310
Armillaria mellea, culture studies, 482
wilt-inducing pigment from, 486
ARMSTRONG, G. M., 465, (465)
ARMSTRONG, J. K., (465), 465
ARNDT, C. H., 220
ARNY, D. C., (391), 465
Arthrobotrys sp., nematode capture by, 405
Ascochyta abelmoschi, in culture, 293
on lupine, 294
phaseolorum and *gossypii* compared, host range, 469
temperature, optimum for growth, 294
gossypii, *abelmoschi* and *phaseolorum* compared, host range, 469
cotton dusting for control, 292
temperature, optimum for growth, 294
imperfecta, alfalfa, effect on seed crop in field and greenhouse tests, 378
fungicide control tests, 379
host range, 378
phaseolorum, *abelmoschi* and *gossypii* compared, host range, 469
spp., cross-inoculations, okra, bean and cotton, 469
ASHRAF, MUHAMMAD, (580)
Aspen, white heartrot, 699
Aspergillus clavatus, antibiotic production in soil and manures, 108
niger, on soybean, 292
spp., on peanuts in market, 474
ATKINSON, D., (587)
ATP, 224
Aureogenus magnivena, *Agalliopsis novella*, transovarian passage in, 9
vastans, *Agallia constricta*, transovarian passage in, 10
Aureomycin, see under Antibiotics
Autogenous necrosis, cabbage, 415
Autoradiography, detection and assay of radioactive toxins *in vivo*, 237
Avena barbata, 310
Avocado, Phytophthora root rot resistance, 490
AYCOCK, ROBERT, 50
Azalea, dagger nematode, 296

Bacillus phytophthorus, effect on respiration of potato, 222
polymyxa, effect of, on tissue culture, 92
spp., in seeds of certified beans, 483
pepper fruit spot disease in Palestine, 92
soft rot in various vegetables, 92
subtilis, antibiotic, toximycin from, 23
effect of, on tissue culture, 92
Bacitracin, see under Antibiotics
Bacteria, ring-rot on potato bags, ultra-violet, X-ray and nuclear radiation, effect of, on, 440
Bacterial blight, chrysanthemum, 522
soybean, inoculum dilution and host reaction, 468
pathogen in seed and soil, 189
canker, tomato control, 473
leaf spot pathogens association with soybean seed, soil and debris, 189
metabolites, effect of on carrot tissue, 92
pustule, soybean, pathogen in seed and soil, 189
spot, broccoli, 583
spring canker, peach seasonal occurrence, 465
stem rot, geranium, 588
wilt, carnation, cause of wilting, 587
carnation, virulence, retention in culture, 406
tobacco, 240
Bacteriophage, *Xanthomonas pruni*, factors affecting plaque formation, 27
BAINES, R. C., 465
BAKER, KENNETH F., (470), 585
BALD, J. G., 141, 146, 151, 167, 585, (587)
Barberry, Japanese, cucumber mosaic virus host, 489
Barley, cereal yellow-dwarf virus, aphid vectors, 309, 312
epiphytotic conditions favoring, 310
effect of air temperature and nitrogen supply on injury by olefinic peroxide fumigants, 588
ergot associated with loose smut, 461
Helminthosporium spp., reactions to isolates of, 517
powdery mildew, environmental conditioning of plants, effects on susceptibility to, 162

Barley—*continued*
Pythium graminicola, varieties resistant to, 474
seedling blight control, 391
Septoria leaf blotch, host-parasite relationships, 473
smut, covered, control, 393
inoculating method, 697
loose, control, 393, 313
ergot associated with, 461
races, 407, 482
resistance of crosses to races of, 371
viability of seeds of parental varieties and crosses inoculated with pathogen, 370
varieties, differential test plants for loose smut, 407
wheat striate mosaic virus on, 539
yellow-dwarf virus, 128
effect on yield, 134
Barnett, H. L., 341
Barrus, M. F., 350
Bart, George J., 466
Basidiobolus spp., odor of benzene hexachloride and of *Streptomyces* spp. produced by, 405
Basidiomycete, alfalfa winter crown rot, HCN production by, 478
host parasite relation, 478
species, hydrogen cyanide production, alfalfa crown rot, and effect of temperature and media, 581
Bassett, E. G., (110)
Baxter, John W., 178
Baxter, L. W., 466
Bayberry-yellows virus, transmission, 15
Bean, alfalfa black stem on, 378
mosaic strains on, 271
yellow mosaic virus on, 38
anthracnose, pressure effects on, 70
Ascochyta spp. on, 469
broad, cucumber mosaic virus strains on, 334
clover-root nematode on, 606
cucumber mosaic virus strains on, 334
halo blight, control, streptomycin, 407, 480
lima, Cercospora purple stain, seed, 477
cucumber mosaic virus strains on, 334
downy mildew, 419
potassium nutrition, relation to root-knot nematode development in, 171
root-knot nematode on, 171
microorganisms in certified seeds of, 483
Mung, alfalfa black stem on, 378
olefinic peroxide leaf injury and protectants, 588
powdery mildew, control in laboratory, 477
pressure effects on, 70
rice, alfalfa black stem on, 378
cucumber mosaic virus strains on, 334
root rot control, soil fumigation, 590
rust, pressure effects on, 70
seed, orange color caused by *Coynebacterium flaccumfaciens*, strain of, 483
snap, Cercospora purple stain, seed, 477
southern mosaic virus, serological test for, 77
sting nematode, 295
streptomycin movement in, 480
Tepary, cucumber mosaic virus strains on, 334
tobacco mosaic virus, acquired resistance, 490
necrosis virus strain, adapted to, 590
velvet, alfalfa black stem on, 378
white mold, occurrence in Idaho, 519
yellow mosaic virus, age of plant and method of inoculation, 473
necrotic-lesion producing strain (new) on, 45
strains on, 271
Beckman, C. H., 441, 448
Beet, seedling necrosis, fungicide in soil and fertilizer tests, 590
stubby-root nematode, 296
Belonolaimus gracilis, see also Nematodes, sting
control, 291, 438
on various crops, 291, 434
2-benzothiazolylthioglycolate, sodium and potassium salts of, formative effects on tomato, effects on chemical composition on tomato plants, in chemotherapy of Fusarium wilt, 43
Beraha, Louis, 383, 466
Berbee, Flora, (466)
Berbee, J. G., (448), 466
Berberis Thunbergii, see Barberry, Japanese, 489
Bergeson, G. B., (264)
Berkeley, G. H., 111
Beta vulgaris, see also Beet, bayberry yellows virus, 18
Bever, Wayne M., 681
Biographical sketch, Forest Milo Blodgett, 350
E. W. Bodine, 60
Luther Parris Byars, 59
Melville Thurston Cook, 591
Richard English Vaughan, 349
James Stewart Wiant, 592
Biological control, possible explanation of, 701
Birch yellow, canker and decay, *Poria laevigata*, incitant, 487
Birchfield, Wray, 289
Birdsfoot trefoil, *Stemophylium* loti sp. nov., 577
Black, L. M., 9, (387), 466
Black medic, alfalfa black stem on, 378
red clover vein-mosaic virus, 473
Black root rot, tobacco resistance to, 239
Black rot, apple, effect of continual use of fungicides, 268
sweet potato, differences in susceptibility of varieties to, 78
Black shank, tobacco, resistance to, 239
Bloch, Robert, 351
Blodgett, Forest Milo, Biographical sketch, 350
Bloom, James R., 466
Blueberry, blossom and twig blight control, 481
stem canker, resistant varieties, 589
Bodine, E. W., Biographical sketch, 60
Bolton, A. T., (345)
Bonde, Reiner, 463
Boosalis, M. G., (401), 467
Boothroyd, C. W., 467
Borlaug, Norman E., 467, (468)
Botryodiplodia phaseoli, *Macrophomina phaseoli* as synonym, 610
Botryosphaeria ribis, apple rot in Maryland, 407
muscadine grape, 629
shell bark relation to, 467
Botrytis cinerea, blueberry blossom and twig blight control, 481
methyl bromide control, 376
potato, effect of maturity and water content of tubers on susceptibility to, 338, 339
effect of peptic enzymes on, 340
shell bark relation to, 467
strawberry gray mold control, 482
gladiolorum, control, 151, 154
corm lesions, 146
heat-curing of corms infected by, 141
neck rot humidity, effect on, 169
host tissue defensive reactions, 170
leaf spot, comparison with, 167
origin of, 167
soil temperature, effect on, 169
sporulation, 169
spp., onion blast incitants, 483
Bouchea prismatica, spot anthracnose, 406
Bouchereau, Paul E., 289
Bouteloua cupitpendula, 310
Boyce, John S., Jr., 644
Boyle, John S., 467
Boyle, Lytton W., 571
Brakke, Myron K., 387, 467

BRANDT, WILLIAM H., 467
BRANN, J. L., (480)
Brassica juncea, B. nigra virus property studies, 541
green peach aphid virus transmission, 209
nigra, virus, aphid-host plant interaction variation, 209
aphid transmission, 209, 254
host range, 541
properties: thermal inactivation, dilution, end-point, longevity *in vitro*, 541
oleracea, see Cabbage
BRAUN, ARMIN C., 204
BRENER, W. H., (466)
BRIERLEY, PHILIP, 404 (406)
Broccoli bacterial spot, *Pseudomonas maculicola*, incitant, 583
Bromus spp., 310
Brown patch, susceptibility of various lawn grasses to, 110
Brussels sprouts, *Heterodera cruciferae* on, 260
Buchloe dactyloides. see under Grasses, buffalo
Buffer systems, tomato early blight control and fungus inhibition, 478
Buprestidae, see under Insects
BURKHOLDER, W. H., (350), 522
Butterfly pea, alfalfa black stem on, 378
Byars, Luther Parris, Biographical sketch, 59

Cabbage, autogenous necrosis, inheritance of, 415
Badger Market, new disease resistant variety, 649
black ring virus, effect of air temperatures on, in *Nicotiana* spp. 550
Fusarium, metabolites production of vascular browning, 490
stubby-root nematode, 296
virus A, effect of air and soil temperature on, in *Nicotiana* spp., 550
yellows, 278, 650
resistance, basis of, 489
Caeoma spp., nomenclature, changes in, 183
CAIRNS, ELDON J., 105, 404
Cajanus indicus, see Pigeon pea
CALAVAN, E. C., 467
Calyptospora columnaris, 360
Camellia, spp., *Glomerella cingulata* on varieties of, 466
leaf and flower variegation, graft transmitted, 293
CAMPANA, RICHARD J., 468
CAMPOS, T. A., 468
Cantaloupe, cucumber Fusarium on, 481
Capsicum, see under Pepper
Captan, see also under Fungicides, 58
Carbon^{14}, see Radioisotopes
Carnation, bacterial wilt, cause of wilting, 587
clover-root nematode on, 606
fumigant injury, 489
Fusarium stem rot, 485
greasy blotch, 585
Carneocephala flaviceps, see Insects, yellow-headed sharpshooter
CARPENTER, T. R., 156
Carrot, bacterial metabolites, effect of on tissue culture, 92
tissue culture, effect of bacteria on, 92
CARTER, J. C., 468
Caryophyllaceae, carnation Fusarium wilt, susceptibility to, 465
Castanea spp., see Chestnut
CATION, DONALD, 468, (471)
Celery, cucumber mosaic virus strains on, 336
disease resistance, breeding for, 254
seedbed disease control, 293
sting nematode, 293, 295
stubby-root nematode, 296
western mosaic, aphid transmission, 29
CENOZ, HUGO P., (487)
Cerambycidae, see under Insects
Ceratocystis fimbriata, black cane rot of *Syngonium auritum*, form of, 586
Cercospora apii, celery, resistance, 254
beticola, sugar beet breeding for resistance to, 298
seminalis, on buffalo grass, etiology of, 488
spp, on soybean, 292
soybean seeds, purple strain, 477
Cereal yellow-dwarf virus, see under Viruses
Cerosporella spp., nomenclature, changes in, 183
CERVANTES, JAVIER, (480)
CETAS, R. C., (467)
Chaetomium spp., in soil, inhibitory to damping-off of peppers, 586
Chalara fagacearum, association with other wood rotting fungi, 467
oak wood, effect on, 467
quercina, see also *Endoconidiophora fagacearum* carbon, sources for growth in culture, 445
heterothallism, 67
insects associated with, 69
nitrogen sources for growth in culture, 444
oxidizing and reducing agents effect on growth in culture, 442
pathogenicity, 68
perithecial stage of, 65
pH effect on growth in culture, 442
vitamins and purines for growth in culture, 443
CHAMBERLAIN, D. W., 468
CHANDLER, W. A., 108
Charcoal rot, effect of temperature and moisture on growth, 634
linear growth rates of the fungus, 633
pycnidial stage of fungus, 608
Chayotes, stubby-root nematode, 296
Chemotherapeutants, benzothiazole, 44
2-benzothiazolythioglycolic acid, derivatives, Dutch elm disease, 486
potassium and sodium salts of, comparison of fungitoxic and chemotherapeutic potency, 44
calcium sulfamate, pH effect on activity, wheat rusts, 498
wheat leaf and stem rust control, 497
4-carbomethylaminophenyl sodium sulfonate, wheat leaf and stem rust control and phytotoxicity, 497
2,4-dichlorophenoxyacetic acid, Fusarium wilt of tomato, effect upon, 137
metabolism of host, effect upon, 139
3,5-dinitrobenzoic acid, wheat stem rust, 661
ferbam, tobacco blue mold, 486
hydrazines, wheat leaf and stem rust control and phytotoxicity, 497
indole-3-acetic acid, Fusarium wilt of tomato, effect upon, 137
metabolism of host, effect upon, 139
mercaptobenzothiazole, 44
α-naphthaleneacetic acid, Fusarium wilt of tomato, effect upon, 137
metabolism of host, effect upon, 139
α-naphthoxyacetic acid, Fusarium wilt of tomato, effect upon, 137
metabolism of host, effect upon, 139
nitrated and hydroxy benzine derivatives, wheat stem rust, 662
oxyquinoline benzoate and sulfate, oak wilt, 475
phenylhydrazine, pH effect on activity, wheat rusts, 498
wheat leaf and stem rust, control and phytotoxicity, 497
picric acid, wheat stem rust, 661
potassium α(2-benzothiazolyl) thiopropionate, potassium β(2-benzothiazolyl) thiopropionate, 44
2-thiazolylthioglycolate, 44
8-quinolinol, effect on Fusarium wilt of tomato, 283
effect on toxicity of lycomarasmin, 284
8-quinolino sulfate, Rhizoctonia, 475
sulfamates, wheat leaf and stem rust, control and phytotoxicity, 496

Chemotherapeutants—*continued*
sulfanilic acid, wheat leaf and stem rust control and phytotoxicity, 497
thiazole, 44
2,3,5-triiodobenzoic acid, Fusarium wilt of tomato, effect upon, 137
metabolism of host, effect upon, 139
2,4,6-trinitrobenzoic acid, wheat stem rust, 661
zineb, tobacco blue mold, 486
Chemotherapy, Dutch elm disease, 486
Fusarium wilt of tomato, effect upon, 137
oak wilt, 475
plant growth-regulators, 137
radiophosphorus inhibition of tobacco mosaic virus, 208
tobacco blue mold control with zineb and ferbam, 486
wheat rusts, 496, 659
Chenopodium album, tobacco-etch virus, local lesion host, 568
Cheo, Pen Ching, 78
Cherewick, W. J., 461, (697)
Cherry, disease control, mist and hydraulic spraying compared, 465
flower inoculum for virues, 479
fluoride concentrations, foliar, factors affecting, 587
green ring mottle virus, 481
quantitative method of assay on cucumber, 176
leaf spot, control, 109, 468
recurrent chlorotic ring pattern effect of aging on infectivity of isolates in inoculum from vegetable marrow and pumpkin, 325
ring spot virus, effect on growth of nursery trees, 480
Sclerontinia fructicola, and *S. laxa*, pathogenicity compared, 589
tatter leaf virus, infectivity and longevity, effect of temperature and age of source of inoculum on, 326
quantitative method of assay on cucumber, 176
trunk and crown rot, British Columbia, 402
virus complex, flower inoculum for isolations, 479
virus seed transmission, quick method, 467
yellows virus, infectivity of isolates, 325
quantitative method of assay on cucumber, 176
Chestnut blight, tannin factor in resistance, 32
Chick-pea, blight fungus, breeding for resistance, 580
Childs, James F. L., 99, 101
China aster, effect of methyl bromide on, 376
Chinn, S. H. F., 701
Chloramphenicol, see under Antibiotics
Chloris gayana, 310
Chlorogenus myricae, 19
p-chlorophenyl p-chlorobenzene sulfure, see Insecticides, miticides, Ovotran
Chloromycetin, see under Antibiotics
Chloropicrin, see under Soil fumigants
Christensen, J. J., 468, (479)
Christiansen, D. W., (483)
Christie, J. R., 295
Chromatography, detection and assay of radioactive toxins *in vivo*, 237
Chrysanthemum, bacterial blight, dissemination and varietal susceptibility, 524
pathogen: *Erwinia chrysanthemi*, 526
symptoms, 522
methyl bromide injury, 489
stunt virus, 406
Marmor chrysanthemi, proposed name, 406
tomato aspermy virus on, 404
Cicer arietinum, see Chick-pea
Cinnamon, C. A., (439)
Cintractia spp., nomenclature, changes in, 183
Cirsium arvense, *Verticillium albo-atrum* on, 588
Citrus, concentric canker, 99
feeder roots, microorganisms isolated from, 395
Fomes applanatus associated with wood rot on, 99
Citrus—*continued*
Fusarium spp. on feeder roots, 395
gummosis, 101
paradisi, see Grapefruit
reticulata, see Tangerine
sinensis, see Orange
tristeza, evidence of in Florida, 51
grafting experiments, 51
wood rot of, 99
Cladosporium cucumerinum, cucumber breeding for resistance to, 257
cucumber varieties, new, resistant to, 215
effect on tissues of resistant and susceptible cucumber varieties, 481
fulvum, tomato varieties immune from, 109
spp. on soybean, 292
Clarkia elegans, *Phytophthora parasitica*, susceptibility to infection by, 290
Clavacin, see Antibiotics, Patulin
Clayton, E. E., 239, 468
Clitoria mariana, see Butterfly pea
Clover, alfalfa black stem on, 378
alsike, clover-root nematode on, 606
red clover vein-mosaic virus, 473
crimson, cucumber mosaic virus strains on, 336
for bioassay of wound-tumor virus, 387
Ladino, alfalfa mosaic strains on, 271
clover-root nematode on, 606
mosaic virus complex, 271
effect on yield, 273
red clover vein-mosaic virus, 473
yellow bean mosaic strains on, 271
red, bean yellow mosaic virus, 470
vein-mosaic, age of plant and method of inoculation, 473
incidence and host hange, 473
sweet, necrotic-lesion strain of yellow bean mosaic virus on, 46
Pythium spp., pathogenicity on seedlings, 474
white, clover-root nematodes on, 603
sweet, cucumber mosaic virus strains on, 336
red clover vein-mosaic virus, 473
wound-tumor virus, insect vectors, 9
yellow sweet, red clover vein-mosaic virus, 473
Coccomyces hiemalis, see Cherry leaf spot
control by actidione, 468
Cochliobolus sativus, inheritance of characters, 487
Cochran, L. C., 585, (691)
Colladonus geminatus, see Insects, under leafhoppers,
Colletotrichum destructivum, grass and legume isolates compared, 487
on brome grass, 486
gossypii, survival on cotton seeds in storage, 220
hibisci, kenaf varieties resistant to, 647
lagenarium, pH effect on lesions on cucumber leaves, 489
watermelon breeding for resistance to, 254
lindemutheanum, pressure effects on, 70
phomoides, perfect stage, 285
Common vetch, alfalfa black stem on, 378
Conifer, damping-off, seedlings, control, 446, 466
seedlings, disease complex etiology and control, 469
Conks, sterile of *Fomes igniarius*, 699
Conover, Robert A., 344
Cook, Harold T., 592
Cook, Melville Thurston, Biographical sketch, 591
Cooley, J. S., (50)
Coons, George H., 297, 405
Cooper, D. C., (490)
Cooper, W. E., 289
Corbett, M. K., 469
Corn, awl nematode, 296
charcoal rot, pycnidial stage of fungus on kernels, 608
cucumber mosaic virus strain on, 336
Gibberella stalk rot, loss of leaf area effect, 477

Corn—*continued*
Hormodendrum kernel rot, 386
injury, autogenus, 386
internal temperatures of root and stem compared with soil, 317
Pythium root rot, effect of aldehyde on virulence of pathogen, 289
spp. on seedlings, effect of temperature and moisture, 477
seedling blight control, 477
seedling root rot, 476
smut, gall development, effect of temperature on, 656
partial vacuum method inoculating seedlings, 654
stalk rot effect on yield, 479
sting nematode, 295, 435
stubby-root nematode, 296
Corticium galactinum, white pine needle blight studies, 468
praticola, as perfect stage of *Rhizoctonia solani*, 488
Corynebacterium flaccumfaciens, causing orange color in bean seed, 483
michiganense, tomato seed transmission, 473
sepedonicum, potato bags, ultra-violet, X-ray and nuclear radiation, effect of, on, 439
potato resistance to, 251
Cotton, *Ascochyta* blight, dusting for control, 292
gossypii on, 294
spp. on, 469
fiber properties and wilt resistance, new hybrids, 292
Fusarium wilt, metabolites production of vascular browning, 490
new resistant hybrids, 292
resistance, basis of, 489
nematode-wilt complex, soil fumigation for, 292
Rhizoctonia solani, dusting for control, 292
root-knot on Deltapine 15, 292
root rot, tests for control, 589
sting nematode, 434, 438
causing, loss of resistance to Fusarium wilt, 475
thrips, dusting for control, 292
Verticillium wilt, 585
breeding for resistance to, 489
varieties reaction to, 489
COURTNEY, W. D., (598)
Cowpea, alfalfa black stem on, 378
alfalfa yellow mosaic on, 39
Cercospora purple stain, seed, 477
cucumber mosaic virus strains on, 334
Fusarium wilt, races on, 291
necrotic-lesion strain of yellow bean mosaic virus on, 46
sting nematode, 435
COX, R. S., (419), 469
Criconema spp., see Nematodes, ring
Criconemoides spp., see Nematodes, ring
CRITTENDEN, H. W., 405
Cronartium quercuum, pycniospore stage on *Pinus taeda*, 488
strobilinum, pycniospore stage on *Pinus caribaea*, 488
CROSSAN, D. F., 469
Crown gall, organization of tumor cells in presence of naphthalene acetic acid, 204
Crucifer viruses, thermal inactivation and dilution end-point of strains, 545
Cucumber, anthracnose, effect of pH on development of lesions on leaves, 489
downy mildew, breeding for resistance to, 257
Fusarium wilt, 481
mosaic virus, breeding for resistance to, 256
powdery mildew, breeding for resistance to, 257
Rhizoctonia foliage blight on, 290
scab, breeding for resistance to, 257
effect of pathogen on tissues of resistant and susceptible varieties, 481
resistant varieties, new, 215
sour cherry tatter leaf virus isolates in, 326
Cucumber—*continued*
stone-fruit, green ring mottle virus isolates in, 325
latent virus transfer to, by cherry flower inoculum, 479
necrotic ringspot virus isolates in, 325
viruses, test plant for virus assay, 175
tomato aspermy virus on, 404
Cucumis sativus, see Cucumber
Cucurbits, disease resistance, breeding for, 254
inhibition of stone-fruit virus by diluents of, 328
stone-fruit virus isolates in, 324
Cultures, preservation of under oil, 407
CUMMINS, GEORGE B., (178)
CUNNINGHAM, H. S., 95
CURL, E. A., 61, (287), 469
Curled dock, clover-root nematode on, 606
Curvularia lunata, "fading-out" on turf grass, control, 109
spp. on soybean, 292
Cuscuta spp., bayberry-yellows virus, 15
Cyamopsis tetragonoloba, see Guar
Cylindrocladium scoparium, control on conifer seedlings, 469
Cycloheximide, see Antibiotics, actidione
Cynodon dactylon, 310
Cynosurus echinatus, 310

Dactylella sp., nematode capture by, 405
Dactylis glomerata, 309
Damping-off, conifer seedlings, control, 469
control by copper sulphate, 125
sugar beet complex, 300
temperatures of root and stem effect on, 317
tobacco seedlings, Venezuela, 125
DARLEY, ELLIS F., 469, (588)
DARLING, H. M., (476)
Dasyscycha willkommii, on larch, disease survey, 405
Datura spp., tomato bushy-stunt virus strains, test plant, for 485
stramonium, alfalfa yellow mosaic on, 39
cucumber mosaic virus strains on, 334
Daucus carota, bayberry-yellows virus on, 18
DAVIES, M. E., (109)
DAVIS, DAVID, (43), 137, 470
DAVIS, LILY H., (470), (585), 586
DEAN, JACK L., 290
DELP, CHARLES J., (475)
Dendrophoma obscurans, on strawberry, control, 471
DESJARDINS, P. R., 687
DEVAY, J. E., 470, (654)
DIACHUN, STEPHEN, 470
Diaporthe citri, shell bark relation to, 467
Diazouracil, effect on multiplication of TMV, 589
2,3, dichloro-1,4-naphthoquinone, see Fungicides, Phygon
2,4-dichlorophenoxyacetic acid, Fusarium wilt of tomato, effect upon, 137
metabolism of host, effect upon, 139
dichloropropene-dichloropropane, see Soil fumigants, DD
DICKSON, JAMES G., 349, (473), (478), (527), (581)
DIENER, T. O., 586
3,5-dimethyltetrahydro-1,3,5,2-thiadiazine-2-thione, see Fungicides, Crag 974
DIMOCK, A. W., (522)
DIMOND, A. E., 43, (137), 195, 229, (281), 319, 470, (488), 619, 663
Dinitrocarprylphenyl crotonate, see Fungicides, Karathane
2,4 dinitrophenol, effect on plant respiration, 224
Diplocarpon earliana, strawberry, photographic infection scales for roguing susceptible seedlings, 345
Diplodia natalensis, on peanuts in market, 474
shell bark relation to, 467
zeae, effect on corn yield, 479
Disease development,
Gladiolus corms, effect of heat-curing upon, 141
effect of humidity upon, 142

Disease resistance, alfalfa to leaf spot, method of measurement, 651
avocado to Phytophthora root rot, 490
barley varietal crosses to loose smut races, 369
bean to TMV, acquired, 490
cabbage, Badger Market, new variety to yellows and mosaic, 650
to yellows, basis of, 489
celery, breeding for, 254
chick-pea to blight fungus, breeding, 580
cotton and cabbage vascular Fusaria effect of culture filtrates on, 489
to Fusarium wilt, 292, 489
to Verticillium wilt, breeding, 489
cucumber to scab, host cell reaction, 481
cucurbits, breeding for, 254
flax to rust, 624
varieties to Fusarium wilt races of pathogen, 495
identifying method, interspecific transfer, 468
kenaf to *Colletotrichum hibisci*, 647
multiple, in cotton, 292
in oats, breeding, 291
in sugar beets, breeding, 295
oats to crown rust, effect of temperature and age of plant, 484
Persea spp. to Phytophthora root rot, 490
plant growth-regulators inducing, 137
potato, nature of, 245
varieties, *Phytophthora infestans* races in Mexico, 454
sugar beet to curly-top virus strain, 588
varieties and inbreds resistant to virus yellows, 405
tobacco breeding for, 469
black shank field inoculation effectiveness for testing resistance, 289
tobacco, nature of, 243
to black shank altered by root-knot nematode, 483
to mosaic virus, 475
tomato, to *Cladosporium fulvum*, 109
Fusarium wilt resistance affected by metabolic inhibitors, 483
wheat breeding, rust, "composite variety," 467
rust differences in seedling and adult stage, 468
stem rust breeding, 472
linkage between resistance to race 15B and susceptibility to races 49 and 139, 472
to bunt, conditions affecting, 473
Dithane D 14, see Fungicides, Nabam
Dithane Z 78, see Fungicides, Zineb
Ditylenchus dipsaci, see under Nematodes
sp., culture of, for use in teaching and research, 105
Dolichodorus heterocephalus, see Nematode, awl
Dolichos lablab, alfalfa yellow mosaic on, 39
DOOLITTLE, S. P., 333, (404)
Dothiorella sp., apple, associated with cankered branches, 407
DRECHSLER, CHARLES, 405
Drosophila melanogaster, see under Insects
Dry rot, sprout inhibitors, effect of, on—in stored potatoes, 95
DUNEGAN, J. C., 405
DURBIN, RICHARD D. 470, 586
DURRELL, L. W., 60

EARHART, R. W., 290, 516
EDGERTON, C. W., 591
Eggplant, *Ascochyta* spp. on, 469
cucumber mosaic virus strains on, 334
potato virus Y on, 217
Verticillium albo-atrum isolates on, 466
EIDE, CARL J., (486)
EISENSTARK, A., (27)
Electrophoretic fractionation, tobacco ring spot virus, 687
mobility, tobacco ring spot virus, 690
tobacco mosaic virus, groups of strains, compared, 687
ELLIS, D. E., 290, (293), (469)
Elm, Dutch elm disease therapy of, 486
Elsinoe boucheae, spot anthracnose of *Bouchea*, 406
Elymus canadensis, *Ustilago striiformis* and *U. bullata* hybrids on, 548
spp., 201, 310
EMERSON, F. H., (109)
Empoasca fabae, *Solanum* spp., resistance to, 246
ENCISO, T., (472)
Endive, effect of air temperature and nitrogen supply on injury by olefinic peroxide fumigants, 588
Endoconidiophora fagacearum, see also *Chalara quercina*
chemotherapy, 475
compatibility types A and B from naturally and artificially inoculated red oaks, 341
compatibility types in trees, 645
distribution in trees, 644
Drosophila melanogaster vector, 466
effect of temperature and humidity on longevity of spores, 476
fungus pads, 287
growth and development on C^{14}-labeled medium, 490
longevity in oak lumber and products, 485
Nitidulidae associated with, 466
fimbriata, differences in susceptibility of sweet potato varieties to, 78
Endothia parasitica, tannin factor in resistance to, 32
Endothlaspis spp., nomenclature, changes in, 183
Endria inimica, see Insects, under leafhoppers
ENGLISH, HARLEY, 586
Entyloma, nomenclature, changes in, 183
Enzyme action, control of in cell, 221
effect of quinones and ketones on, 226
β-glucosidase, Fusarium wilt infected tomatoes, 470
pectic, of *Botrytis cinerea* on potato tissue, effect, 340
of *Pythium debaryanum* on potato tissue, effect, 340
role in Fusarium wilt of tomato, 535
pectin methyl esterase, tomato Fusarium wilt, 472, 535
pectinase, *Chalara quercina*, produced in culture of, 447
pectinesterase, vascular browning, 490
polygalacturonase, tomato Fusarium wilt, role, 535
vascular browning, 490
polyphenol oxidase, *Sclerotinia fructicola*, relation to fungitoxicity, 483
Stemphylium sarcinaeforme, relation to fungitoxicity, 482
respiration, effect of toxins on, 221
tomato reaction and enzyme activity, 536
vivotoxins, 232
Eragrostis cilianensis, wheat striate mosaic virus on, 539
Ergot, loose smut associated with, 461
ERWIN, DONALD C., 586
Erwinia amylovora, greenhouse test for removing pear seedlings susceptible to infection by, 161
pear, inoculation of seedlings, 161
susceptibility tests, 405
Pyrus spp. crosses, effect upon, 161
atroseptica, black leg on potato, caused by, 92
chrysanthemi, bacterial blight of chrysanthemum, 526
physiological studies, 525
phytophthora, 92
spp., chrysanthemum, cross-inoculation tests, 525
effect of, on culture tissue, 92
cultural characters of, 526
Erysibe spp., nomenclature, changes in, 184
Erysiphe cichoracearum, culture on sunflower tumor tissue *in vitro*, 343
graminis, effect on respiration of wheat leaves, 223
environmental conditioning, effects on susceptibility of barley, 162
tritici, temperature effect on, on wheat seedling varieties in greenhouse, 290
polygoni, bean, control in laboratory, 477
effect on respiration of clover leaf, 223
pressure effects on, 70

Ethyl alcohol, pea, triple response to, 664
tomato Fusarium wilt, 663
Ethylene, auxin-ethylene balance, 232
effect on respiration of apples, 225
effect on respiration of lemon and pear, 222
epinasty in Fusarium wilt of tomato, 470, 664
pea, effect on, 664
production by *Fusarium oxysporum* f. *lycopersici*, 664
toxic action, mode of, 232
Ethyl mercuric p-toluene sulfonanilide, see Fungicides, Ceresan M
Ethyl mercury chloride, see Fungicides, Ceresan

False smut, on buffalo grass, etiology, 488
FAVRET, EWALD A., 471
FEAZELL, GEORGE D., 290
FEDER, WILLIAM A., 471
FELDMESSER, JULIUS, (471), 471
FELIX, E. L., 290
FERGUS, CHARLES L., 103
FERGUSON, JOHN, (375), 471, 586, (593)
Ferric dimethyl dithiocarbamate, see Fungicides, ferbam
Festuca spp., 309, 310
Figs, Kadota, Alternaria rot control, 586
Cladosporium surface mold control, 586
Fir, Douglas, disease complex control, 469
FISCHER, GEORGE W., 181, (200), 547
FISHER, H. H., (45), (407)
Flax, Fusarium wilt races, varietal reaction to, 493
rust, epidemiology in North Central States, 624
physiologic race survey from 1931, 625
races, genetic characters, 626
varieties resistant to, fungus races interaction, 625
Septoria linicola, dissemination, 468
FLETCHER, D. G., 471
FLOR, H. H., 624
Fluoride, concentrations in foliage, factors affecting, 587
FOLSOM, DONALD, (108)
Fomes applanatus, concentric canker, associated with, 99
wood rot of citrus, associated with, 99
igniarius, aspen, white heartrot, 699
sterile conks on aspen, 699
FOSTER, H. H., 290
FOWLER, DORSEY L., (277)
FOWLER, MARVIN E., 405, 406
FRENCH, ALEX M., 513
FRIEDMAN, BERNARD A., 108, 412
FROEHLICH, H. P., (477)
FULTON, ROBERT H., 471
FULTON, ROBERT W., 472
Fumigation, peroxide, olefinic bean leaf injury and protectants, 588
effect of air temperature and nitrogen supply on injury to herbaceous plants, 588
Fungi, effect of different pathogens on temperature of infected leaves, 679
mycorrhiza-like fungus in potatoes, 490
physiology, oxidizing and reducing agents effect on *Chalara quercina* growth in culture, 442
sclerotia, viability vs. colonization by soil organisms, 471
sterile conks, *Fomes igniarius*, 699
wood rotting in buildings, 20
Fungicidal action, nature of, 476
Fungicides, general, combined soil and foliage treatment, conifer seedling complex, 469
effect of fertilizers on effectiveness in soil, 590
liquid concentrates vs. dusts for onion smut control, 480
methods for evaluating effects on *Verticillium* in soil, 593
mixtures in control of apple scab and quince rust, 109
polyelectrolytes as spray conditioners, 489
sensitivity to, difference in single conidial isolates, 490
soybean autumn seed treatments, 476
spore germination and hyphal growth, differential action, 476
Fungicides, general—*continued*
testing of fungicides on apple seedlings in greenhouse, 109
tests for fungicidal solution strength for bulb dips, 598
uptake from dilute solution by fungus spores, 480
vapor action, mercury seed treatments, oats, 465
Fungicides, Aagrano 350, peanut seed treatment, 291
aconitic acid, fungicidal action, 476
Agrox, oat seedling blight, vapor effect, 465
aniline derivatives, fungicidal action, 476
Arasan, see also thiram
Arasan SF, soybean seed treatment, 476
biphenyl derivatives effect on apple fruit flavor, 385
effect on *Monilinia fructicola* on apples, 383
bis(p-Nitrophenol) carbonate, peach brown rot, 290
Bordeaux, actidione compared, cherry leaf spot, 468
apple black rot, 268
conifer seedling disease complex, 469
effect of continual use on black rot of apple, 268
effect of spraying muscadine grapes, 630
injury, effect on temperature of leaves, 678
oak leaf-blister, 472
Cadsol, 109
calcium polysulphides, root rot, cotton and alfalfa, tests, 589
Calocure, conifer needle injury, 469
captan (N-trichloromethylthio tetrahydrophthalimide), 58
and actidione mixture, wheat covered smut, 461
apple scab and quince rust, 109
bean halo blight, 407
blueberry blossom and twig blight, 481
oak leaf-blister, 472
peach brown rot, 407
peach canker, 109
peanut seed treatment, 291
rose blackspot, 406
strawberry gray mold, 482
strawberry leaf blight, 471
tomato damping-off, 291
C.C.C. 224, and 5400, peanut seed treatment, 291
Ceresan, (ethyl mercury chloride) peanut seed treatment, 291
shallot pink root, 294
test for solution strength, bulb dip, 602
Ceresan M (ethyl mercuric p-toluene sulfonanilide) oat seedling blight, 465
seed treatment, barley smut, 393
barley seedling blight, 392
oat, 290
oat, Helminthosporium blight, 391
oat smut, 393
wheat bunt, 393
wheat loose smut, 315
sodium nitrate effect on effects of, in soil, beet seedling necrosis, 590
test for solution strength, bulb dip, 598
Ceresan, New Improved (ethyl mercury phosphate) ferbam compared with, in control of corm-borne disease of gladiolus, 154
gladiolus corm-borne disease control, 154
Lysol compared with, in control of corm-borne disease of gladiolus, 154
root injury to gladiolus, 154
soybean seed treatment, 476
test for solution strength bulb dip, 602
tomato damping-off, Vancide 51 interaction, 291
chloranil (tetrachloro-para-benzoquinone) alfalfa black stem, 379
barley loose smut seed treatment, 313
gladiolus corms, 429
peanut seed treatment, 291
shallot pink root, 294
soybean seed treatment, 476

Fungicides—*continued*

chlorinated hydrocarbons, root rot, cotton and alfalfa, control tests, 589
o-chlorobiphenyl, effect on *Monilinia fructicola*, 383
Cop-O-Zink, oak leaf-blister control, 472
copper, impregnated paper wrap in peach rhizopus and brown rot, 407
strawberry leaf blight control, 471
Copper A, mist and hydraulic spraying, 465
copper oxide, conifer needle injury, 469
copper oxysulfate in fungiside-insecticide mixtures, roses, blackspot, 406
copper sulphate, damping-off in tobacco seedbeds, 125
tobacco seedlings, effect on growth, toxicity, 125
copper, tri-basic, polyelectrolytes as spray conditioners, 489
sulphate, oak leaf-blister, 472
zinc and DDT combination, cotton dusting, 292
Crag 341 (2-heptadecyl glyoxalidine acetate) apple scab and quince rust, 109
effect on fig, Cladosporium surface mold and Alternaria rot, 586
Crag 974 (3,5-dimethyltetrahydro-1,3,5,2-thiadiazine-2-thione) *Fusarium moniliforme*, 465
nematodes, 465
Penicillium sp., 465
Pythium spp., 465
Cuprocide, tomato damping-off, Vancide 51 interaction, 291
dimethyldithiocarbamic acid, sodium salts of and 2-mercaptobenzothiazole mixture, conifer damping-off soil treatment, 466
Dow-9-B, gladiolus corm injury, 429
Dowicide A, peach brown and rhizopus rots, 407
Elgetol (sodium salt of dinitro ortho cresol), Tag 331 compared, strawberry leaf blight, 472
ethylenebisthiuram sulfides, bean powdery mildew, 477
ferbam, (ferric dimethyl dithiocarbamate) apple, black rot, 268
apple scab and quince rust, 109
bean halo blight, 407
blueberry blossom and twig blight, 481
conifer seedling disease complex, 469
gladiolus corm-borne disease, 151
peach canker, 109
strawberry leaf blight, 471
tobacco blue mold, 486
tomato injury, 291
tomato, damping-off, 291
p-fluorobiphenyl, effect on *Monilinia fructicola*, 383
formaldehyde, conifer seedling disease complex, 469
damping-off, conifer seedlings, 469
root-rot, conifer seedlings, 469
tests for solution strength, bulb dip, 598
fungicide 406, see captan
heavy metals, fungicidal action, 476
2-heptadecyl-2-imidazaline, uptake from solution by fungus spores, 480
heterocyclic compounds, fungicidal action, 476
hydrocarbons, fungicidal action, 476
insecticide, dust mixtures, rose blackspot, 406
Karathane (dinitrocarprylphenyl crotonate) bean powdery mildew, 477
lime-sulfur, apple black rot, 268
injury, effect on temperature of leaves, 678
Lysol, gladiolus corm-borne diseases, 151
magnetic sulfur, apple black rot, 268
Manzate (manganese ethylene bisdithiocarbamate) apple black rot, 268
blueberry blossom and twig blight, 481
combined with insecticides, rose blackrot, and effect on plant, 406
conifer seedling disease complex, 469
fig, Cladosporium surface mold and Alternaria rot control by and phytotoxicity, 586
nitrate fertilizer effect on effects of in soil, 590
oak leaf-blister, 472
peach brown rot, 290
peach, effect of color on, 290
protectant against olefinic peroxide injury to beans, 588
Mathieson 275, peanut seed treatment, 291
2-mercaptobenzothiazole and dimethyldithiocarbamic acid mixture, conifer damping-off soil treatment, 466
Merculine, oak leaf-blister, 472
mercuric chloride, shallot pink root, 294
Mersolite 8 (phenyl mercuric acetate), test for solution strength, bulb dip, 598
Microgel, oak leaf-blister, 472
Monsanto 4367, 4370, oak wilt, 475
nabam and zinc sulphate, oak leaf-blister, 472
naphthoquinones, chromatographic separation of, 110
effect upon respiration, 226
effect upon sulfhydryl containing enzymes, 226
orthocide, see Captain, coined name, 58
Panogen (methyl mercuric dicyan diamide) oat seedling blight, 465
oat seed treatment, 290
test for solution strength, bulb dip, 598
Penn Salt NP770, peach brown rot, 290
phenols, fungicidal action, 476
N-phenylmercuriethylenediamine, apple scab and quince rust, 109
phenyl-mercury compounds, bluegrass, *Helminthosporium vagans*, 476
Phygon (2,3,dichloro-1,4-naphthoquinone) alfalfa black stem, 379
apple scab and quince rust, 109
blueberry blossom and twig blight, 481
peach canker, 109
shallot pink root, 294
Phygon XL, effect on fig, Cladosporium surface mold and Alternaria rot, 586
oak leaf-blister, 472
PMAS, conifer needle injury, 469
polyethyl biphenyl, effect on *Monilinia fructicola*, 383
Puratized, conifer needle injury, 469
shallot pink root, 294
quaternary ammoniums, fungicidal action, 476
quinones, fungicidal action, 476
naturally occurring, 226
Semesan, alfalfa black stem, 379
shallot pink root, 294
setrete, oat seedling blight, vapor effect, 465
Spergon, see chloranil
SR-406, captan, coined name, 58
sulfur, ferbam, insecticide mixture, rose blackspot and effect on plants, 406
magnetic "70," apple scab and quince rust, 109
wettable, bean leaf injury, 477
bean powdery mildew, 477
mist and hydraulic spraying, 465
peach brown rot, 290
with BHC and DDT, cotton dusting, 292
Tag 331 (phenyl mercury acetate), 109
strawberry leaf blight, 471
strawberry leaf injury, 472
Tersan, alfalfa black stem, 379
tetrachloroethylene, peach rhizopus rot fumigation, 407
tetramethylthiuram sulfides, bean powdery mildew, 477
thiocarbamates, fungicidal action, 476
thiram, (tetramethylthiuram disulfide) alfalfa black stem, 379
bean powdery mildew, 477
conifer damping-off soil treatments, 466
conifer seedling disease complex, 469
corn seedling blight, 477

Fungicides, thiram—*continued*
onion smut seed treatment, 596
peanut seed treatment, 291
pelleting conifer seed, 466
shallot pink root, 294
soybean seed, 189, 476
tomato damping-off, Fermate interaction, 291
Thiuram, peach canker, 109
toximycin as, 23
trichloroethylene, peach rhizopus rot fumigation, 407
undecylenic acid, oak wilt, 475
Vancide 51 (sodium salts of dimethyl dithiocarbamic acid and of 2-mercapbenzothiazole)
blueberry blossom and twig blight, 481
fig Cladosporium surface mold and Alternaria rot, 586
oak leaf-blister, 472
oak wilt, 475
peach brown rot, and plant injury, 290
tomato damping-off, Ceresan, New Improved, and Cuprocide interaction, 291
Vancide 51 ZW, soybean seed treatment, 476
Zinc Coposil, fig Cladosporium surface mold and Alternaria rot, 586
zinc sulphate, tri-basic copper combination, cotton dusting, 292
zineb, (zinc ethylenebisdithiocarbamate) conifer seedling disease complex, 469
effects of spraying compared with Bordeaux, muscadine grape, 631
fig Cladosporium surface mold and Alternaria rot, 586
oak leaf-blister, 472
protectant against olefinic peroxide injury to beans, 588
tobacco blue mold, 486
ziram (zinc dimethyl dithiocarbamate) blueberry blossom and twig blight, 481
fig Cladosporium surface mold and Alternaria rot, 586
Fungitoxicity, benzothiazole and related compounds to Fusarium wilt, 43
Sclerotinia,—of benzothiazole and related compounds to, 43
Stemphylium,—of benzothiazole and related compounds to, 43
Fungus sectoring, in *Fusarium lini*, 488
Fungus species, on peanuts in market, 474
Fungus spores, fungicides, uptake by, 480
Fusarium bulbigenum, 397
conglutinans, races 3 and 4, 465
races causing wilt of cowpea, 291
Elegans section, citrus feeder roots, 395
equisetti, citrus feeder roots, 395
lateritium f. *pini*, host range and geographical distribution, 475
lini, chromatic variants in, 488
Martiella section, citrus feeder roots, 395
moniliforme control, 465
survival on cotton seeds in storage, 220
oxysporum, 397
on soybean, 292
tobacco varieties resistant to, 240
toxic metabolites, 116, 490
f. *apii*, 254
f. *callistephi*, methyl bromide control, 376
f. *conglutinans*, cabbage resistance to, 649
races 1, 2, and 3, 280
f. *cubense*, soil moisture and type, effect on growth and survival, 499
corm lesions, 146
control, 151, 154, 376
heat-curing of corms infected by, 141, 425
"walling-off" by cork tissue, 430
wound periderm formation in corms related to, 425
f. *lini*, flax varieties reaction to races, 493
methods of inoculation, 491
pathogenic clones, selection and maintenance, 492
Fusarium—*continued*
races, pathogenicity, 493
soil type and temperature effect on virulence, 492
f. *lycopersici*, ammonium-N and nitrate-N levels in culture filtrate, 196
chemotherapeutic activity of benzothiazole and related compounds, 43
ethanol and ethylene production, 470, 664
β-glucosidase production of, 470
lycomarasmin, 116, 195
magnesium sulfate effect of, 466
nutritive sprays on host effect on, 466
osmotic properties of leaf parenchyma in diseased plants, 619
pectic enzymes, 472, 535
physiology, 195
polysaccharides in culture filtrates, 196
resistance affected by metabolic inhibitors, 483
sensitivity of single conidial isolates to fungicides, 490
toxin demonstrated by intergeneric grafts, 470
transpiration, losses and velocity, 619
vascular reaction of host in tomato to, 116
f. *melonis*, muskmelon resistance to infection by, 255
f. *niveum*, cucumber Fusarium cross-inoculations, 481
watermelon resistance to, 255
f. *pisi*, races 1 and 2, resistant pea variety, New Era, 473
f. *raphani*, 277
roseum, carnation clones identical with *F. roseum* f. *cerealis*, 485
cross-pathogenicity tests, 485
sambucinum, 95
semitectum, citrus feeder roots, 395
solani, effect on respiration of potato tubers, 222
var. *Martii*, 397
sp., on cucumber, 481
wound cork, effect of on penetration of, 95
spp., celery seedbed, control, 293
isolation from plant tissues, improved method, 700
tobacco seed-bed incidence, 125
vasinfectum, nematode complex on cotton, soil fumigation, 293
soil microorganisms growth in presence of, 482
Fusicoccum persicae, control, 109
FUTRELL, MAURICE C., 290, 472

GAINOR, G., 509
GALLICCHIO, VIRGINIA, (478)
Galls, *Agrobacterium tumefaciens*, colonial variants, stability in, 510
cedar-apple rust in tissue culture, 360
organization and recovery of crown-gall cells in presence of α-naphthalene acetic acid, 204
tissue in gummosis of grapefruit, 101
tomato, golden nematode, cytology, 471
GARREN, KENNETH H., (644)
Gastridium ventricosum, 310
Genetic factors, chick-pea resistance to blight fungus, 580
in *Cochliobolus sativus*, 487
color and mating type in *Helminthosporium sativum*, 487
disease resistance, method of identifying suited to interspecific transfer, 469
in flax to rust, 626
in wheat, resistance to yellow rust, 471
Genetics of microorganisms, *Glomerella cingulata*, linkage groups, 489
graminicolus smut fungi, in induced hybridization, 547
Melampsora lini, genes for virulence, 626
Ustilago striiformis hordei and *U. bullata* hybrids, inheritance of sorus and spore characters, 547
zeae, pathogenicity of biotypes, method to determine, 654
variability in sensitivity to fungicides, 490

Geranium, bacterial stem rot, 588
GERDEMANN, J. W., 603
Gibberella saubinettii, effect on respiration of potato slices, 222
zeae, corn stalk rot, effect of loss of leaf area, 477
effect on corn yield, 479
GIBLER, J. W., (468), 472
GIDDINGS, N. J., 587, (684)
GILMER, R. M., 108
GINOZA, W., 587
Gladiolus, bean virus 2 affecting, 111
bean yellow mosaic virus, 114
Botrytis neck rot, 167
cork formation, conditions favoring, temperature optimum, 426
corm-borne disease control, 151
corm lesion incidence, 146
corm mummifiaction, 430
corms, *Botrytis* on, 141, 146
disease control by heat-curing, 141
Fusarium oxysporum on, 141, 146
Penicillium spp. on, 149
Rhizoctonia solani on, 148
Rhizoglyphus rhizophagus, on, 150
Sclerotinia on, 148
wound periderm, effect of heat-curing upon, 141
cucumber mosaic virus affecting, 111
flower inoculum for virus isolations, 479
fungicide injury, 429
Fusarium basal rot, 146
yellows, heat treatment for control, 425
"walling-off" by cork tissue, 430
wound periderm formation related to, 425
histological studies of tissue reaction to Botrytis neck rot, 170
root injury, ethyl mercury phosphate, 154
Stemphylium leaf spot, *Pleospora* sp., conidial stage, 585
tobacco ringspot virus affecting, 112
virus complex, 114
Globe amaranth, cucumber mosaic virus strains on, 336
Glomerella cingulata, antibiotic assay, for, 23
camellia dieback and canker, 466
genetics of mutant characters, linkage groups, 489
muscadine grape, 629
glycines, 487
Glomerella phomoides Swank sp. nov., 285
Glucose-1-phosphate, effect on multiplication of TMV, 589
Glycine max, see Soybean
GNAGY, R. M., (587)
Gnomonia veneta, on white oak, weather related to occurence of, 103
Gomphrena globosa, see Globe amaranth
GOLD, A. H., (206), 458
GOLDEN, A. MORGAN, 406
GOODE, MONROE J., 472
Gossypium spp., Verticillium wilt, reaction to and breeding for resistance, 489
GOTHOSKAR, S. S., 472, (483), 535
GOTO, S., 472
GOTTLIEB, DAVID, (478)
GOULD, C. J., (598)
Grafting, citrus triteza studies, 51
transmission of phony disease virus, 692
GRAHAM, J. H., 189, 193, 577, 642
GRAHAM, S. O., 587
GRAHAM, T. W., 291, (291), 434, (475)
Gram, see Chick-pea
Gramicidin, see Antibiotics
Graminicolus smut fungi, induced hybridization, 547
GRANT, THEODORE J., 51
Grape, degeneration related to Pierce's disease, 611
Macrophoma ripe rot, 629
Melanconium leaf and stem fleck on, 347
muscadine, angular leaf spot, 629
effect of spraying on yields and disease incidence, 629
Grape—*continued*
rots: black and bitter, 629
shelling of berries, varietal variation, 632
Pierce's disease virus, insect vectors, 293, 612
yellow mosaic virus, 475
Grapefruit, actinomycete associated with gummosis of, 101
wood rot and canker associated with *Fomes applanatus*, 99
Grass, blue, control of diseases on varieties, 476
brome, *Colletotrichum destructivum* on, 486
brown patch, susceptibility of various lawn grasses to, 110
buffalo, false smut, etiology, 488
cereal yellow-dwarf virus, immune species, symptomless carriers, 310
Curvularia "fading-out," 109
head smut of, differentiation of races, 201
orchard, purple leafspot, effect of periods of high and low humidities on, 642
seed treatment, 487
striped smuts, host range and distribution, 527
GRAVES, CLINTON H., JR., 473
GREEN, G. J., 473
GREEN, MAURICE, 473
GREENLEAF, W. H., 564
GRIES, GEORGE A., (343)
GRIFFITH, R. B., 473
GRISWOLD, CHARLES L., (466)
GROGAN, R. G., 473
GROSSBARD, ERNA, 108
Growth regulators, see Plant growth regulators
Guar, alfalfa black stem on, 378
Guayule, charcoal rot, 633
GUBA, E. F., 109
Guignardia bidwellii f. *muscadinii*, 629
Gummosis, actinomycete, associated with on grapefruit, 101
GUNKEL, W. W., (480)
Gymnoconia peckiana, 360
Gymnosporangium juniperi-virginianae, mycelium on callus in tissue culture, 362
sporulation on galls in tissue culture, 362

HAASIS, F. A., (469)
HAFIZ, ABDUL, 580
HAGEDORN, D. J., 473
HALLER, MARK H., (406)
HALPIN, J. E., 474
HAMILTON, J. M., 109
HAMMACK, MARIAN G., 587
HANSON, E. W., (473), (474)
HARDING, R. B., (477)
HARE, W. W., 291, 474
HARRISON, A. L., 291
HART, HELEN, (485)
HASKETT, W. C., 474
HAYDEN, E. B., 474
HEIBERG, BARBARA C., 474
HEIM, JEAN M., 343
Heliconia spp., resistance to *Marasmius stenophyllus*, 479
Helicotylenchus spp., see Nematodes, spiral
Helixin B, see Antibiotics
Helminthosporium, inoculum preparation technique, 401
sativum, inheritance of characters, 487
spores in soil, method of estimating population, 478
wheat crown and root rot, 484
spp., control, 391
differential host reactions and conidial measurements, 516
vagans, bluegrass, control, 476
victoriae, 230
oat, breeding for resistance, 291
vapor action of mercury seed treatments, oats, 465
HEMERICK, G. A., (378)
HENDERSON, L., (468)
HENDRIX, J. W., 587, (643)

HENRY, A. W., 461
HENRY, BERCH W., 81, 474
HENSON, LAWRENCE, (470)
2-heptadecyl glyoxalidine acetate, see Fungicides, Crag 341
HEPTING, GEORGE H., 475
HESS, G. E., (687)
Heterodera cruciferae, on Brussels sprouts, 260
rostochiensis, see also Nematode, Golden and tobacco cyst nematode, host preferences, 106
schachtii var. *trifolii*, see also Nematodes, clover-root life history, pathogenicity and host range, 603
sp., host preference, 106
HEWITT, WM. B., 475
Hibiscus cannabinus, see Kenaf
HICKMAN, DONALD, (486)
HILBORN, M. T., 475, (481)
HILDEBRANDT, A. C., (92), 475, (490)
HOFFMAN, P. F., 475, (490)
HOLDEMAN, Q. L., (291), 291, (434), 475
HOLLIS, JOHN P., 355
Hollyhock, *Ascochyta* spp. on, 469
HOLMES, FRANCIS O., 475
HOLTON, C. S., 219, 322, 398, (558)
HOLTZMANN, O. V., 587
HOOKER, ARTHUR L., 476
HOPPE, PAUL E., 386
Hopperburn, potato, resistance to, 252
Hordeum nodosum, 201
spp., 310
Hormodendrum corn kernel rot, 386
HORNER, C. E., 587, 588
Horseradish mosaic virus, effect of air temperatures on, in *Nicotiana glutinosa*, 550
white rust, overwintering, 485
HORSFALL, JAMES G., 476, (482)
HOTSON, HUGH HOWISON, 360, 659
HOUSTON, BRYON R., (128), 271, (309), (458), 491
HOWARD, F. L., 109, 476, (486)
HUBERT, ERNEST E., 403, 637
HUGHES, M. B., (50)
Humidity, Botrytis neck rot of gladiolus, relation to, 167
effect on longevity of spores of oak wilt pathogen, 476
effect on production and survival of sporangia of *Phytophthora infestans*, 507
gladiolus corms, effect on disease development on, 142
HUNGERFORD, C. W., 519
HUTCHINS, LEE M., 691
Hydnaceae, wood rotting by 21
Hydrangea macrophylla, tomato ringspot virus on, 404
Hydrogen cyanide production by alfalfa crown rot pathogen, 581
Hydrogen-ion concentration, antibiotic production in soil and manures effect on, 108
Chalara quercina, effect on growth, 442
Colletotrichum lagenarium, effect on lesions on cucumber leaves, 489
Fusarium oxysporum f. *lycopersici* culture filtrates, 195
wheat leaf and stem rust, effect on chemotherapeutic action, 498
HYRE, R. A., 419

Indole acetic acid, see Plant growth regulators
Inheritance of characters, *Glomerella cingulata*, linkage groups, 489
graminicolus smut fungi, in induced hybridization, 547
Inheritance of resistance, barley to loose smut races, 369
cabbage to yellows and mosaic, 650
chick-pea to blight fungus, 586
flax to rust, 624
kenaf to *Colletotrichum hibisci*, 647
multiple, sugar beet, 297
oats to cold, 291
tobacco to mosaic virus, 475
wheat crosses to yellow rust, 471
Inheritance of susceptibility, to autogenous necrosis, cabbage, 415
Inoculum, flower tissue, cherry, virus complex, 479
flower tissue, gladiolus, cucumber mosaic virus, 479
leaf and flower compared, stonefruit viruses, 480
plant disease fungi, large quantity preparation, 401
Insecticides, BHC, cotton, 292
DDT, cotton, 292
lindane-fungicide mixtures, roses, effect on, 406
wheat stem rust, 661
lindane and DDT-fungicide mixtures, roses, effect on, 406
miticides, Aramite in fungicide-insecticide mixtures, roses, effect on, 406
Ovotran, fungicide-insecticide mixtures, roses, effect on, 406
Systox, *Ditylenchus dipsaci* feeding habits related to effect on, 471
Rhobditis sp. feeding habits related to effect of, 471
Insects, *Aceria tulipae*, wheat streak virus, survival at low temperatures, 484
wind dispersal, 485
Agallia constricta, New Jersey potato yellow-dwarf virus, 10
wound-tumor virus, 387
Agalliopsis novella, wound-tumor virus, 9, 387
Agrilus bilineatus associated with oak wilt, 61
aphids, apple-grain, (*Rhopalosiphum prunifoliae*) cereal yellow-dwarf virus, 309
cereal yellow-dwarf virus, 128, 312
corn (*Rhopalosiphum maidis*) cereal yellow-dwarf virus, 132
grass (*Mocrosiphum dirhodum*), cereal yellow-dwarf virus, 132
false cabbage (*Rhopalosiphum pseudobrassicae*) Brassica nigra virus, 542
greenbug, cereal yellow-dwarf virus, 309
green peach aphid (*Myzus persicae*), Brassica nigra virus, 209, 542
potato virus A strains, 479
potato virus Y, 217
tomato aspermy virus, 404
English grain (*Macrosiphum dirhodum*), cereal yellow-dwarf virus, 312
Macrosiphoniella sanborni, tomato aspermy virus, 404
Myzus solani, tomato aspermy virus, 404
pea aphid (*Illinoia pisi*), Wisconsin pea streak virus, 484
Rhopalosiphum rufomaculatum, tomato aspermy virus, 404
potato virus Y, mechanical inoculation vs. aphid inoculation, 469
western celery mosaic, 29
Buprestidae associated with oak wilt, 62
Cerambycidae associated with oak wilt, 61
Drosophila melanogaster, oak wilt, 466
Hessian fly, wheat, crown and foot rot relation to, 467
hopperburn, potato resistance to, 252
leafhopper, *Carneocephala flaviceps*, grape Pierce's disease virus, 293
Colladonus germinatus, peach yellow leaf roll virus transmission studies, 561
Endria inimica, wheat striate mosaic virus, 537
injection techniques, 388
Solanum spp. resistance to, 245
for bioassay of wound-tumor virus, 382
Nitidulidae, oak wilt associated with, 456
oak wilt associated with, 61, 69
Rhizoglyphus rhizophagus, gladiolus corms, 150
root mites on gladiolus corms, 150
Scolytidae associated with oak wilt, 61
Septoria linicola dissemination, 468
thrips, cotton dusting for control, 292
transovarian passage of New Jersey potato yellow-dwarf virus, 10
of wound-tumor virus, 9

Insects—*continued*
viruses, loss of tranmissibility by, 466
yellow-headed sharpshooter Pierce's disease, 612
Internal cork disease, sweet potato, 50
Ipomoea batatus, see Sweet potato
Isoelectric point, tobacco mosaic virus, groups of strains compared, 587
Isolation of fungi from woody tissue, instruments, 403
Isoleucine solutions effect on multiplication of TMV, 589
Itersonilia sp., parsnip, manner of invasion, 485
IVANOFF, S. S., 291

JEDLINSKI, HENRY, (467)
JEFFERS, W. F., (406)
Jelly end rot, potato, 467
JENKINS, W. R., (406)
JENSEN, D. D., 561
JEWELL, FREDERICK F., 476
JOHNSON, H. W., 476, (477)
JOHNSON, S. P., 363
JONES, E. D., 476
JONES, FRED REUEL, 651
JONES, L. S., (585)
Juniperus virginiana, callus tissue in culture, *Gymnosporangium* on, 360
Jute, charcoal rot, pycnidial stage of fungus on, 608

KAESBERG, PAUL, (482)
KEIL, H. L., 477
KEITT, G. W., (23), 59, (391), (478), (481)
KELLER, JOHN R., 406
KELMAN, ARTHUR, 291
Kenaf, *Colletotrichum hibisci*, resistance to, 647
KENDRICK, E. L., 477
KENDRICK, JAMES B., SR., (473)
KENDRICK, J. B., JR., 477, 588, (588)
KERNKAMP, M. F., 378, 477, (584)
Ketones, effect upon respiration, 226
unsaturated, naturally occurring, 226
KEVORKIAN, ARTHUR G., 406
KILPATRICK, R. A., 292, (476), 477
KING, J. F., (487)
KLOS, E. J., (481)
KNOWLES, P. F., (491)
KOEHLER, BENJAMIN, 477
KOMMEDAHL, THOR, 478
KONZAK, C. F., 369
KOPPE, R. K., (587)
KUNTZ, J. E., (441), (448)

LACKEY, C. F., 588
LAMBERT, E. B., (471)
LANGLEY, B. C., (291)
Larch, European, canker disease survey, 405
LARSON, R. H., (479), 596, (649)
Lathyrus odoratus, see Sweet pea
LEANDER, F. JOHNSON, (479)
LEBEAU, J. B., 478, 581
LEBEN, CURT, (23), 391, (465), 478
LEDINGHAM, R. J., 478, (701)
LEGATOR, MARVIN, 478
Legumes, Cercospora purple stain on seeds, 477
seed treatments, 487
LEHMAN, S. G., 292, 460
Lemon, shell bark, fungi and wound healing relation to, 467
Leptographium sp., distribution, and inoculation studies, 637
on pine, western white and lodgepole, 637
symptoms, histology and phytotoxin tests, 639
Leptothyrium pomi, 471
Lespedeza, clover root nematode on, 605
Rhizoctonia aerial blight, soil source of inoculum, 293
Lethum australiense, see also Viruses, 467
Lettuce, Romaine, effect of air temperature and nitrogen supply on injury by olefinic peroxide fumigants, 588
Light effect on lesion formation in potato virus Y on *Physalis floridana*, 2
effect on reaction of Kubanka wheat to races of *Puccinia graminis tritici*, 481
effect on susceptibility of oat hybrids to stem rust, race 7, 473
effect on thiouracil inhibition of TMV, 555
effect on thiouracil up-take by tobacco leaves, 556
LINFORD, M. B., (603)
LINN, M. B., (466), (485)
LIVINGSTON, J. E., 496
LOCKWOOD, J. L., 478
LOEGERING, WILLIAM Q., 479
Loganberry, root rot, *Phytophthora* sp., *Pythium* sp., and nematode associated with, 588
Lolium multiflorum, 310
Lonicera hispidula, apple flyspeck host, 586
Lotus spp., *Stemphylium loti* on, 579
LOWNSBERY, B. F., 106
LUCAS, G. B., (289), (483), (701)
LUKE, HERBERT H., 479
Lupine, *Ascochyta* spp. on, 294
Cercospora purple stain, seed, 477
LUTTRELL, E. S., 347, 629
Lycium spp., potato virus A strains on 479
Lycomarasmin, characteristics of, 231
chelation with metals, 233, 282, 320
complexes, iron and copper relative stability of, 284
decomposition to non-toxic substance I, 319
decomposition rate *in vitro*, 320
effects on transpiration, compared with *Fusarium oxysporum* f. *lycopersici*, 116
ferro-lycomarasmin, decomposition compared with, 320
inhibition by copper ion, 282
inhibition by iron chelating agent, 281
inhibitors as chemotherapeutants, 283
iron complex, 116
non-competitive inhibitor, 233
physiology of production in cultures of *Fusarium oxysporum* f. *lycopersici*, 195
potentiation by iron ion, 281
production rate estimates *in vitro*, 320
8-quinolinol, effect on, 283
role in Fusarium wilt of tomato, 284
strepogenin, inhibited by, 233
temperature and pH effect on life of, 320
toxicity, 116, 199, 281
Lycopersicon esculentum, see Tomato
peruvianum, 109
root-knot nematode resistance, 290
pimpinellifolium, potato virus A strains on, 479
spp., buckeye rot, susceptibility to, 290

MACDONALD, H. A., (487)
MACLACHLAN, D. S., 479
Macrophomina phaseoli, as synonym for *Botryodiploida phaseoli*, 610
on peanuts in market, 474
pycnidial stage, 608
Macrosiphum spp., cereal yellow-dwarf virus vector, 132
Magnamycin, see Antibiotics
Malas baccata, *Venturia inaequalis*, races 2 and 3 on, 484
Maleic hydrazide, increase in size of pustules of wheat leaf rust, 497
MANDELL, J. D., 27
Manganese ethylene bisdithiocarbamate, see Fungicides, Manzate
MANKIN, C. J., (587)
MARAMOROSCH, KARL, (387)
Marasmius stenophyllus, sheath rot, abaca, 479
Marigold, tissue culture, effect of bacteria on, 92
Marmor chrysanthemi, see Chrysanthemum stunt virus
cucumeris, muskmelon, breeding for resistance, 256
medicaginis H. var. **flavovarians** N. var., 38
upsilon, see Viruses, potato Y

MARSHALL, BARTON H., JR., 425
MARTIN, J. P., (465)
MARTIN, W. J., 292, (292), 432
MAUGHAN, F. B., (477)
MAYHEW, D. J., (587)
MCBETH, C. W., 264
MCCALLAN, S. E. A., D1, (480)
MCFADDEN, LORNE A., (522)
MCINTOSH, D. L., 402
MCKEEN, W. E., 588
MCWHORTER, FRANK P., 479, (479)
MEINERS, JACK P., 200
Melampsora helioscopiae, effect on respiration of *Euphorbia peplus* leaves, 223
lini, 360
genetics of pathogenicity, interaction of races and flax varieties, 626
physiologic specialization of races, 624
survey of pathogenicity and epidemiology in North Central States since 1931, 624
Melanconium fuligeneum, grape, leaf and stem fleck on, 347
muscadine grape, 629
Melanin, amino acid stimulation of, in *Streptomyces*, 357
ascorbic acid in relation to, 361
formation in galls of cedar apple rust, 361
production correlated with growth of *Streptomyces*, 358
Melilotus alba, see Clover, white sweet
Meloidogyne arenaria, on pepper varieties, 474
hapla, on pepper varieties, 474
incognita, bean, 171
ovicidal and therapeutic value of various compounds, 485
potassium nutrients of host, effect upon development of, 171
sweet potato resistance to, 290
tomato, 171, 290
var. *acrita*, 292, 474
spp., see also Nematodes, root-knot, 260, 292
cotton, soil fumigation, 293
Fusarium wilt complex on cotton, 293
MENZIES, J. D., 684
Metabolism, effect of toxins on, 221
polyphenol oxidase system in cedar-aple rust fungus, 361
Methionine, effect on toxins, 233
Methocel, in seed treatment, 476
Methods, see Techniques
Methyl bromide, see Soil fumigants
Methylene blue, wheat stem rust, 661
Methyl mercuric dicyan diamide, see Fungicides, Panogen
MICHAELSON, MERLE E., 479
Microflora, on tobacco roots, diseased, 590
Microorganisms, culture under oil seal, 407
Microthyriella rubi, apple flyspeck, 470
ecology and host range, apple, 586
imperfect stage of, 470
MIDDLETON, JOHN T., (588), 588
MILBRATH, J. A., 479
MILLER, J. B., (489)
MILLER, JULIAN H., 292
MILLER, LAWRENCE P., 480
MILLER, PATRICK M., 480
MILLER, R. L., (461)
MILLER, V. L., (598)
Millet, sting nematode, 295
MILLIKAN, D. F., 480
MILLS, W. R., 454, (456)
MILSTREY, R. A., 406
Mineral element deficiency, see also Nutrition, potassium nutrition of host effect upon development of root-knot nematode, 171
Mint, rust, 178
wild species susceptible to races of mint rust, 178
MISHRA, J. N., 338
MITCHELL, J. W., (407), 480
Monilinia fructicola, biphenyl derivatives toxic to, 385
control with fungicides, peach, 290, 407
sensitivity of single conidial isolates to fungicides, 490
MOON, H. H., (405)
MOORE, M. B., (401)
MOORHEAD, ELLEN L., 73
Muhlenbergia asperifolia, *Tilletia asperifolia* on, spore germination, 219
MUNGER, H. M., 254
MUNNECKE, DONALD E., 375, 588
MURAKISHI, H., (643)
MURPHY, D. R., (109)
MURPHY, H. C., (474)
MURPHY, M. M., (629)
MURPHY, R. P., (487)
Musa spp., *Marasmius stenophyllus* on, 479
Mushroom-spawn nematode, temperature-time gradients in control, 404
Muskmelon, breeding for disease resistance, 255
fruit rots, *Phytophthora* sp. and other fungi causing, 293
weather effects on incidence of fruit rots, 293
Mustard, Brassica nigra virus on species, 543
Mycorrhiza, potato tubers and stems, mycorrhiza-like fungus, 490
Mycosphaerella angulata, muscadine grape, 629
fagariae, strawberry, photographic infection scales for roguing susceptibile seedlings, 345
rabiei, chick-pea breeding for resistance to, 580
Myrica carolinensis, bayberry-yellows virus of, 18
Myrothecium spp. in soil, inhibitory to damping-off of peppers, 586
Myzus persicae, see Insects, aphids, green peach

α-Naphthaleneacetic acid, effect of potato leafroll on response of tomato to, 109
effect on root development by tumor tissue of periwinkle, 204
Fusarium wilt of tomato, effect on, 137
metabolism of host, effect on, 139
methyl ester of, effect of, on Fusarium dry rot in potatoes, 95
1-4 naphthoquinones, see also Fungicides, chromatographic separation of, 110
β-naphthoxyacetic acid, Fusarium wilt of tomato, effect upon, 137
metabolism of host, effect on, 139
Narcissus pseudonarcissus, Ditylenchus-induced leaf "spikkels," cytology and structure, 471
NARVAEZ, I., (472)
Nasturtium mosaic virus, Brazilian, cross-protection tests of strains, 307
distribution of, in inoculated plants, 304
Lycopersicum virus 3 relation to, 306
strains, 307
NATTI, JOHN J., 109
NEAL, D. C., 292
NELSON, PAUL E., 589
NELSON, R. R., (477), 584
Nematocides, 485
assaying techniques, 264
DD, 264
screening, 264
Nematodes, associated with killed sclerotia of *Verticillium*, 594
awl (*Dolichodorus heterocephalus*) control in celery seedbeds, 293
Brussels sprouts, *Heterodera cruciferae* on, 269
capture by, *Arthrobotrys* and *Dactylella* spp., 405
citrus, soil fumigation control, 466
clover-root (*Heterodera schachtii* var. *trifolii*) cyst and larvae measurements, 603
life history and host range, 604
control, Crag 974, 465
crop rotation for control and detection, 260
culture for teaching and research, 105

Nematodes—*continued*

cyst-forming, clover-root, 603
dagger (*Xiphinema americanum*), incidence and host range, 296
detection by fumigation, 260
diagnosis by analysis of mineral elements of plant parts, 260
Ditylenchus dipsaci, Narcissus, leaf "spikkels," cytology and structure, 471
Systox, feeding habits related to action by, 471
water screen testing, 264
sp., laboratory culture of, 105
temperature-time gradients in control in mushroom-spawn, 404
ectoparasitic, incidence and host range, 295
field diagnosis of infestations, 259
field testing, 265
fungi complex control in celery seedbeds, 293
Fusarium wilt complex on cotton, 292
golden (*Heterodera rostochiensis*) tomato, cytological effects of, 471
Helicotylenchus sp. on cotton, 292
laboratory preservation and permanent mounts, 262
meadow (*Pratylenchus*) control, 438
sting-compared, 437
sugarcane, associated with diseased roots of, 289
mushroom-spawn, moisture and heat, effect on control, 404
temperature-time gradients in control, 404
nomenclature problems, 262
Panagrellus redivivus, contact nematicidal value of various compounds, 485
Paratylenchus spp., 297
loganberry root rot, associated with, 588
pine root rot, in relation to, 81
pine seedlings, root rot control, 474
Rhabditis sp., Systox, feeding habits related to action, by, 471
ring, *Criconema* spp. and *Crinonemoides* spp., 297
on peanuts, 297
Procriconema spp., 297
root-knot (*Meloidogyne* spp.), 260
African violet, 406
bean, lima, 171
clean fallow for control on soybeans, 405
complex, control in celery seedbeds, 293
on cotton, soil fumigation, 293
on pepper varieties, 474
on sweet potato, 290
on tomato, 171, 290, 465
ovicidal and therapeutic value of various compounds, 485
potassium nutrients of host, effect upon development of, 171
soil injection screening, 265
tobacco, effect on resistance to black shank, 483
Rotylenchulus reniformis on cotton, 292
soil fumigation effect on population, 474
soil injection screening, 265
soil sampling methods of detection, 261
spiral, on Amaryllis, 297
staining plant tissues for detection, 261
sting (*Belonolaimus gracilis*), body measurements, 436
control, 291, 293, 438
cotton, loss of resistance to Fusarium wilt, 475
cotton wilt complex, 260
incidence and host range, 295, 435
meadow, compared, 437
pathogenicity tests, 436
population trends, effect of various plants in greenhouse pot culture, 291
stubby-root, incidence and host range, 296
sugar-beet, 260
techniques for detecting and investigating, 259
temporary laboratory mounts, 262

Nematodes—*continued*

tobacco cyst, host preference, 106
Trichodorus sp. control in celery seedbeds, 293
Tylenchorynchus sp. on cotton, 292
sugarcane, associated with diseased roots of, 289
Neurospora crassa, 359
NEWHALL, A. G., (254), 480
NEWSOM, L. D., 292
Nicandra physaloides, Phytophthora parasitica, susceptibility, 290
potato virus A strains on, 479
potato virus Y on, 217
NICHOLS, CARL W., 555
Nicotiana glutinosa, horseradish mosaic virus air temperature studies, 550
nasturtium mosaic virus, 306
tomato bushy-stunt virus strains on, 485
rustica, cyst nematode, 106
spp., cabbage black ring and virus A, air temperature studies, 553
cucumber mosaic virus, resistance to, 472
cucumber mosaic virus strains on, 334
disease resistance, 243
potato virus Y on, 217
turnip virus 1, air and soil temperature studies, 550
turnip virus 1, effect of nutrition on multiplication in, 669
tabacum, see Tobacco
NIEDERHAUSER, JOHN S., (454), 456, 480
NIENSTAEDT, HANS, 32
Nitidulidae, 466
Nitrogen, see under Nutrition
Nomenclature, see also Taxonomy
Botryodiploidia phaseoli, Macrophomina phaseoli as synonym, 610
Marmor chrysanthemi, stunt virus, 406
nematodes, 262
species concept for classification of smut fungi, 181
Ustilago striiformis, 530
Non-parasitic disease, cabbage, autogenous necrosis, 415
potato jelly end rot, 467
NORTON, DON C., 633
Nothopanax guilfoylei, see Panax
Nutrient solution, effect on plant regulators, effect on rooting of *Prunus mahaleb*, 587
Nutrition, calcium, effect on tomato and potato susceptibility to *Alternaria solani*, 476
concentration of solutions, effect on growth of *Nicotiana* spp., multiplication of turnip virus 1, 671
magnesium, effect on potato and tomato susceptibility to *Alternaria solani*, 476
nitrogen, effect on multiplication of turnip virus 1 and plant growth, *Nicotiana* spp., 669
effect on potato and tomato susceptibility to *Alternaria solani*, 476
phosphorus and potassium levels in relation to multiplication of TMV, 488
Pseudomonas solanacearum response to sources and osmotic pressure of, 291
supply effect on injury by olefinic peroxide fumigants on herbaceous plants, 588
tomato, TMV concentration in, 477
phosphorus, effect multiplication of turnip virus 1 and plant growth, *Nicotiana* spp., 670
effect on potato and tomato susceptibility to *Alternaria solani*, 476
potassium, bean, effect on root-knot nematode, 171
effect on multiplication of turnip virus 1 and plant growth, *Nicotiana* spp., 670
effect on potato and tomato susceptibility to *Alternaria solani*, 476
of host, effect upon development of root-knot nematode, 171
potato, effect on susceptibility to *Alternaria solani*, 476

Nutrition—*continued*
salicylic aldehyde, effect on virulence of Pythium root rot pathogen, 289
tobacco in relation to multiplication of TMV, 488
mosaic virus effect of nitrate, phosphate and potassium levels on growth of host tissue and virus titer, 475
tomato, effect on susceptibility to *Alternaria solani*, 476
turnip virus 1, effect on multiplication in *Nicotiana* spp., 669
Nutrition of microorganisms, amino acids, *Phytophthora infestans* races A and B response to, 514
ammonium salts, *Phytophthora infestans* races A and B responses, 515
asparagine, *Phytophthora infestans* races A and B responses to, 514
biotin effect on growth of *Chalara quercina*, 443
calcium, inhibition of *Sclerotium rolfsii*, 363
carbon, effect on soil microflora, 365
sources for growth of *Charlara quercina*, 445
Chalara quercina, 441
inositol effect on growth of *Chalara quercina*, 443
nitrogen, amino acids as sources for *Venturia inaequalis*, 481
effect on soil microflora, 365
salts effect on isolates of *Sclerotium rolfsii*, 363
sources for growth of *Chalara quercina*, 444
sources for growth of *Venturia inaequalis*, 481
Pseudomonas solanacearum, nitrogen, effect of osmotic pressure, 291
nitrogen, response to different sources of, 291
Pythium root rot of corn, effect of aldehyde on virulence of pathogen, 289
raw sugar, growth stimulant, *Phytophthora infestans* races A and B, 515
succinic acid, *Phytophthora infestans* races A and B responses, 515
thiamine effect on growth of *Chalara quercina*, 443
vitamins and purines, growth of *Chalara quercina*, 443
yeast extract, growth stimulant, *Phytophthora infestans* races A and B, 515
zinc and silicon or gelatin requirement in rust germ tube development, 483

Oak, laurel, dagger nematode, 296
leaf-blister control, 472
rust, pycniospore stage on pines, 488
white, anthracnose, 103
wilt, carbon sources for growth of pathogen in culture, 445
chemotherapy, 475
compatibility types of fungus in trees, 341, 655
control, 406
distribution of the fungus in trees, 644
effect of temperature and humidity on longevity of spores of pathogen, 476
fungus, 61
fungus pads, 287
host responses, 448
inoculum potential vs. season, 469
insects associated with, 69
insect transmission studies, 466
longevity of pathogen in lumber and wood products, 485
nitrogen sources for growth of pathogen in culture, 444
pathogen on C^{14}-labeled medium, 490
radioactive rubidium in water movement, infected trees, 450
spore germination of pathogen vs. season, 469
status, 406
transpiration of host, 450
tyloses in host tissue, 451
vitamin effect on growth of pathogen in culture, 443
Oats, breeding for resistance to cold, 291
cereal yellow-dwarf virus, 309
aphid vectors, 309, 312
epiphytotic, conditions favoring, 310
crown rust, breeding for resistance, 291
effect of temperature and age of plants on reaction of varieties to, 484
effect of air temperature and nitrogen supply on injury by olefinic peroxide fumigants, 588
Helminthosporium blight, seed treatment, 391
spp., reactions of varieties to isolates of, 517
rust, race 4 of pathogen, rapid spread in Argentina, 487
stem, biotypes within race 7 of pathogen, 472
effect of temperature and light, 472
seedling blight, vapor action of mercury seed treatments, 465
seed treatments, 290, 391, 393
Siberian mosaic virus and wheat striate mosaic compared, 539
smut, control, 393
covered, inoculating method, 697
Victoria blight, breeding for resistance to, 291
wheat striate mosaic virus, 539
yellow-dwarf virus, 130
effect on yield, 135
Ogawa, Joseph M., 589
Okimoto, Marion, (465)
Okra, *Ascochyta gossypii* on, 294
spp., on, 469
Olive, Russian, Phytophthora canker, 468
Onion, "Alternaria blotch," 411
Alternaria spp., pathogenicity on bulbs and leaves, 410
storage, 411
bayberry-yellows virus, 18
blast, *Botrytis* spp., incitant, 483
brown blotch pathogens, 409
leaf spotting or blast, *Botrytis* spp., incitants, 483
purple blotch pathogens, 409
smut, control, thiram seed treatment, 596
liquid concentrates vs. dusts for control, 480
Orange, wood rot and canker associated with *Fomes applanatus*, 99
O'Reilly, H. J., 481
Orellana, Rodrigo G., 125
Orthocide, see Fungicides, captan coined name, 58
Oswald, John W., 128, (271), 309, (458), (473)
Oteifa, Bakir A., 171, (407)
Owen, John H., 481

Palm, Washington, Phytophthora trunk rot, 469
Palmiter, D. H., 109
Panax, ringspot virus, symptoms and transmission, 443
Panicum capillare, 310
Pantothenic acid, 221
Papavizas, George C., 481
Parasites, facultative, and obligate, nature of plant reactions to, 353
relation to host metabolism, 221, 227
Parker, K. G., 481
Parsnip, leafspot and canker, manner of invasion by pathogen, 485
Pasteur effect, effect of antimycin A on, 227
loss of, in wheat infected with powdery mildew, 224
Pate, J. B., 647
Patulin, see Antibiotics
Pea, alfalfa black stem on, 378
clover-root nematode on, 606
cucumber mosaic virus strains on, 334
disease resistant variety, New Era, 473
Fusarium, metabolites production of vascular browning, 490
root rot control, soil fumigation, 590
Wisconsin streak virus, pea aphid vector, 484
Peach, antibiotics, effect on, 405
Armillaria mellea, wilt-inducing pigment from, on, 486
bacterial spot spring canker, 465

Peach—*continued*
brown rot control, 290, 406
canker control, 109
cherry yellows transmission to plum scions, 108
leaf drop control, 405
mosaic virus, spread related to age of infection, 585
necrotic ringspot transmission to plum scions, 108
phony disease virus in tops of trees, 692
rhizopus rot control, 406
Sclerotinia fructicola, and *S. laxa*, pathogenicity compared, 589
western X virus, ninhydrin-reacting substance in leaves, 586
physiological changes induced by, 481
yellow leaf roll virus, leafhopper transmission studies, 561
Peanut, alfalfa black stem on, 378
collar rot, heat canker phase, 571
fungi associated with in market, 474
heat canker, occurrence and symptoms, 571
soil temperature and sky condition effect, 572
nematodes on, 295, 297
Sclerotium rolfsii, pathogenicity of isolates on, 289
seed treatment, 291
Pear, fireblight, age of seedlings, effect on resistance to, 161
crosses resistant to, 156
disease rating, effect of temperature on, 160
inoculation method, seedlings, 156
on seedlings, 405
resistance, growth stage, 157
terminal shoot growth cessation, effect on resistance to, 161
streptomycin, effect on Bartlett seedlings, 405
Pecan, dagger nematode, 296
Pelargonium hortorum, see Geranium
PELLETIER, E. NEIL, 481
PELLETIER, R. L., 481
Penicillin, see Antibiotics
Penicillium digitatum, effect on respiration of lemon, 222
patulum, antibiotic production in soil and manures, 108
sp. control, 465
isolate from soil antagonistic to fungi and bacteria, 479
spp., in soil, inhibitory to damping-off of peppers, 586
on peanuts in market, 474
on soybean, 292
Pepper, *Ascochyta* spp. on, 469
cucumber mosaic virus, alfalfa mosaic virus complex on, 333
damping-off, inhibition by soil organisms, 586
effect of methyl bromide on, 376
nematodes, varieties resistant to, 474
Phytophthora blight, 469
potato virus Y on, 217
tabasco, tobacco-etch virus symptoms on, 565
tobacco-etch virus, multiplication and translocation in, 567
tomato aspermy virus on, 404
Verticillium albo-atrum isolates on, 466
Peppermint rust, method of evaluating severity and control, 587
oil gland destruction and leaf drop, 587
Verticillium albo-atrum, pathogenicity of isolates on, 588
Periwinkle, crown-gall tumor cell organization and recovery in presence of α-naphthalene acetic acid, 204
tissue culture, effect of bacteria on, 92
Peronospora effusa, pressure effects on, 70
manshurica, race 4, 292, 460
schachtii, sugar beet breeding for resistance, 300
tabacina, 239
control, 486, 616
distribution, geographical, 616
overwintering, 616
spore formation and germination, 701
spore showers, 616
PERRY, V. G., 293
Persea spp., Phytophthora root rot resistance, 490
PERSON, L. H., (290), 293, 701
Pestalozzia spp., in soil, inhibitory to damping-off of peppers, 586
PETERSON, E. A., (461)
PETERSON, L. C., (469)
Petunia, cucumber mosaic virus strains on, 336
hybrida, bayberry-yellows virus on, 18
potato virus Y on, 217
tobacco necrosis virus strain on roots, adaptation, 590
tomato aspermy virus on, 404
pH, see Hydrogen-ion concentration
Phalaris spp., 310
Phaseolus spp., see Bean
Phenyl mercury acetate, see Fungicides, Tag 331 and Mersolite 8
Phoma betae, sugar beet seedlings black rot, 300
Phomopsis sojae on soybean, 292
Phony disease, see under Peach
Phyllosticta spp. on soybean, 292
Phymatotrichum omnivorum, cotton and alfalfa root rot control tests, 589
Physalis angulata, tomato aspermy virus on, 404
floridana, potato virus Y, 1, 217
quantitative virus assay, 6
spp., *Phytophthora parasitica*, susceptibility to infection by, 290
tobacco-etch virus on, 568
Physalospora obtusa, apple rot in Maryland, 407
effect of continual use of fungicides, 268
Physiologic specialization, flax rust pathogen races, 624
Fusarium congultinans races 3 and 4, 465
races on cowpea, 291
races 3 and 4, 465
Peronospora manshurica race 4, 292, 460
Puccinia graminis tritici races on Kubanka wheat, 481
menthae, races, 178
Tilletia caries, races, pathogenicity, 398
T. foetida races in mixture, host passage survival studies, 558
Ustilago bullata, 200
nuda races, 369, 407
Venturia inaequalis, races 2 and 3, 484
Physoderma spp., nomenclature, changes in, 183
Phytophago destructor, see Insects, Hessian fly
Phytophthora cactorum, olive, Russian, 468
sweet cherry, 402
capsici, pepper, 469
summer squash, 469
cinnamomi control, 376, 465
root rot of avocado and *Persea* spp., 490
citrophthora, methyl bromide control, 376
cryptogea, alfalfa root rot, 586
fragariae, strawberry red stele soil fumigation and fungicide tests for control, 590
infestans, antibiotics effect on, 463
effect on respiration of potato tubers, 222
phaseoli, growth and viability in culture compared, 423
potato resistant varieties reaction to, 455
races, 480
A and B physiologic differences, 513
in Mexico, 454
survival in mixture, 486
Solanum spp. in Mexico, 456
sporangia production and survival, 505
virulent race of, on resistant tomatoes, 344
parasitica, tobacco breeding for resistance to, 239
toxin production in tobacco and tomato sap, 294
var. *nicotianae*, field inoculation effectiveness for testing resistance, 289
root-knot nematode effect on varietal resistance to, 483
var. *terrestris*, *Clarkia elegans* susceptibility to, 290
immune and resistant plant species, 290
phaseoli, effect of temperature, moisture and medium on growth and viability in culture, 419

Phytophthora phaseoli—*continued*
germination of sporangia and zoospores, 422
homothallism in, 423
infestans compared, 423
production of sporangia, 420
sp., loganberry root rot, associated with, 588
muskmelon fruit rots, 293
palm trunk rot, 470
tobacco black shank relation to, 290
PIERSON, C. F., (215), 481
Pigeon pea, alfalfa black stem on, 378
Pine, lodgepole, pole blight, 637
pitch canker, host range and geographical distribution, 475
pole blight, distribution and inoculation studies, 637
phytotoxin tests, 640
symptoms and histology, 639
root rot seedling control, 81, 474
seedling, nematodes associated with, 474
seedling damping-off, 466
white, blister rust, distribution, effect of microclimate, 487
disease complex control, 469
needle blight, *Corticium galactinum* studies, 468
western, pole blight, 637
Pisum sativum, see Pea
PITTS, RALPH, (519)
PLAKIDAS, A. G., 293, (466)
Plant growth regulators, in chemotherapy, 137
indole acetic acid, effect on Fusarium wilt of tomato, 137
effect on metabolism of host, 139
potato virus Y studies, 469
indolebutyric acid, effect on rooting of *Prunus mahaleb*, nutrient pretreatment effect on effect of, 587
maleic hydrazide, size of wheat rust pustules, 497
α-naphthaleneacetic acid, effect on Fusarium wilt of tomato, 137
effect on metabolism of host, 139
effect of potato leafroll on response of tomato to, 109
effect on root development by tumor tissue of periwinkle, 204
methyl ester of, effect on Fusarium dry rot of potato, 95
β-naphthoxyacetic acid, effect on Fusarium wilt of tomato, 137
effect on metabolism of host, 139
pantothenic acid, 221
Pleospora sp., on gladiolus, 585
Plum, Japanese, cherry yellows, transmission to, 108
necrotic ringspot, transmission to, 108
latent necrotic ringspot, quantitative method of assay on cucumber, 176
myrobalan, necrotic ringspot and cherry yellows, transmission to *Prunus salicina*, 108
phony disease virus in tops of trees, 691
Sclerotinia, fructicola and *S. laxa*, pathogenicity compared, 589
Poa spp., 309
POLLEY, DOROTHY, 598
Polycystis spp., nomenclature, changes in, 183
Polyelectrolytes as conditioners for fixed copper sprays, 489
Polymixin, see Antibiotics
Polyporaceae, wood rotting by, 21
Polyscias guilfoylei, see Panax
POPP, W., 482, 697
Populus tremuloides, see Aspen
Poria laevigata, yellow birch canker and decay incitant, 487
Potato, bacterial wilt, southern, 245
blackleg, 245
black scurf, *Rhizoctonia solani*, in India, 645
Botrytis cinerea, effect of maturity and water content on tubers on susceptibility to, 338, 339
charcoal rot, pycnidial stage of fungus on, 608
curly-top virus strain, 587, 684
disease resistance, 245
Potato—*continued*
diseases, multiple resistance, cultivated and wild species, 246
early blight, 245
effect of pectic enzymes of *Botrytis* and *Pythium* on, 340
Fusarium dry rot, 95
golden and tobacco cyst nematode host preferences, 106
green dwarf caused by strain of curly-top virus, 684
hopperburn, resistance to, 252
insect resistance, wild species, 245
internal brown spot associated with infection by *Verticillium albo-atrum*, 108
jelly end rot, 467
late blight, 245
antibiotics effect on, 463
races of pathogen, 480
resistant varieties reaction to races of in Mexico, 455
sporangia production and survival, 505
survival of races of pathogen in mixture, 486
leaf roll, 109, 245
mosaic virus, alfalfa and yellow bean strains on, 271
mycorrhiza-like fungus in tissue cultures, 490
nutrition effect on susceptibility to *Alternaria solani*, 476
"pink eye," associated with infection by *Verticillium albo-atrum*, 108
Pythium debaryanum, effect of sprouting, pre-heating and water content on susceptibility to, 339
ring rot, 245
on burlap bags, ultra-violet, X-ray and nuclear radiation effect of, on, 440
scab, common, 245
storage behavior when infected with *Verticillium albo-atrum*, 108
Synchytrium endobioticum on wild species in Mexico, 480
tissue culture, effect of bacteria on, 92
tobacco cyst nematode, 106
Verticillium albo-atrum, 108, 245
virus A, 245
strains, 479
strains, *Myzus persicae* vector, 479
virus X, 245
method for identifying immune seedlings, 487
virus Y, 1, 217
aphid transmission, 217
host range in Hawaii, 217
mechanical inoculation vs. aphid inoculation and field reaction, 469
strains of, 5
wart, resistance to, 251
yellow dwarf virus resistance to, 249
POUND, GLENN S., 277, 415, 482, (488), 550, (649), 669
Powdery mildew, barley, environmental conditioning, effects on susceptibility to, 162
cucumber breeding for resistance to, 257
muskmelon, breeding for resistance to, 255
tissue succulence related to susceptibility of barley, 162
POWELL, DWIGHT, (383), (465), 482
POWERS, H. R., JR., (483)
Pratylenchus, see under Nematodes
PRESTON, WILLIAM H., JR., (480)
PRICE, W. C., (73), (509)
Procriconema spp., see Nematodes, ring
Propylene oxide on soil, 108
Protomyces spp. nomenclature, changes in, 183
Prune, dwarf virus, infectivity and longevity of inoculum, 326
fluoride concentrations, foliar, factors affecting, 587
Prunus mahaleb, growth regulators and nutrient pretreatment effect on rooting, 587
salicina, see Plum, Japanese
Pseudomonas caryophylli, cause of wilting, carnation, 587
virulence, retention in culture, 406
fluorescens, *Verticillium albo-atrum* associated with, on potatoes, 108

Pseudomonas—*continued*
glycinea, field and laboratory tests on persistence, soybean, 189
inoculum dilution and soybean reaction, 468
soil and debris, persistence in, 191
lapsa and *P. syringae*, physiological reactions compared, 413
maculicola, broccoli bacterial spot, 583
phaseolicola, control, streptomycin sulfate, 407
solanacearum, nitrogen, response to different sources of in culture, 291
sp., blueberry stem canker incitant, 589
spp., serological tests with, 412
tabaci, field and laboratory tests on persistence, soybean, 189
soil and debris, persistence in, 191
Pseudoperonospora cubensis, muskmelon, breeding for resistance, 255
Pseudopeziza medicaginis, alfalfa resistance, rapid method of measurement of, 651
pure culture maintenance and inoculation procedure, alfalfa, 651
Pseudoplea briosiana, perfect stage of *Stemphylium botryosum*, 477, 584
Puccinia antirrhini, pressure effects on, 70
coronata, oat breeding for resistance, 291
var. *avenae*, germ tube development, zinc and silicon or gelatin requirement, 483
oat varieties, effect of temperature and age of plant on reaction of varieties, to, 484
glumarum, wheat crosses, races on, 471
graminis var. *avenae*, race 4 spread in Argentina, 487
race 7, 472
tritici, breeding for resistance, "composite variety" of wheat, 467
effect of chemotherapeutants on, 496, 661
races, 472, 485
races, Kubanka wheat, effect of temperature and light on reaction to, 481
helianthi, pressure effects on, 70
malvacearum, effect on respiration of *Althea rosea* leaves, 223
menthae, control, 180
epiphytology, 178
physiologic specialization in, 178
rubigo-vera tritici, control by chemotherapeutants, 496
Pyrus spp., fireblight, crosses resistant to, 156
Pythium arrhenomanes control, 465
penicillium antagonistic to, 479
debaryanum, potato, effect of pectic enzymes on, 340
graminicola, barley varieties resistant to, 474
irregulare, conifer damping-off control, 466
sp., loganberry root rot, associated with, 588
spp., celery seedbed, control, 293
corn seed decay and root rot, 476
effect of temperature and moisture on pathogenicity, corn seedlings, 477
pathogenicity on alfalfa, 474
pathogenicity on sweet clover, 474
sugar beet seedlings black rot, 300
ultimum, control, 376, 590
temperature related to damping-off by, 318
Quercus spp., see Oak
Quince Rust, control by fungicide mixtures, 109

RAABE, ROBERT D., 482
Radioactive elements, P^{32} effect on tobacco mosaic virus, 206
P^{32} tobacco plant injury, 207
Radioactive pathogens, 236
Radioactive tracers, C^{14}-labeled sucrose in oak wilt fungus study, 490
Ce^{144}, S^{35} and C^{14} in study of uptake of toxicants by fungus spores, 480
P^{32} tomato Fusarium wilt transpiration studies, 621
rubidium, oak wilt, 448
Radioisotopes, carbon^{14}, effect on microorganisms, 237
for tagging toxins, 236
detection of microbial toxins, 236
tritium, effect on microorganisms, 237
for tagging toxins, 236
Radish, cabbage yellows on, 277
Fusarium oxysporum f. *conglutinans*, 280
Fusarium wilt, cabbage yellows compared, 277
control, 280
races 3 and 4 of pathogen, 465
resistant lines of Scarlet Globe, 280
temperature effect, 277
varietal reaction to, 278
virus 1 group, aphid-transmissable, 545
RAM, C. S. VENKATA, 482, (488)
RAMSEY, G. B., (474), (583)
Ramularia spp., nomenclature, changes in, 183
Rape, Brassica nigra virus on varieties, 543
Raphanus sativus, see Radish
RAPPAPORT, IRVING, 589
RASKI, D. J., 259
RAWLINS, T. E., (89), (206), (589)
RAWLINS, W. A., (350), (480)
RAY, B. ROGER, (486)
RAYCHAUDHURI, S. P., 15
Respiration of pathogen in relation to temperature of host, 679
relation of temperature in diseased, injured and healthy leaves, 679
toxins effect on, in plant tissue, 221
western X virus infected peach vs. healthy, 481
Rhamnus californica, apple flyspeck host, 586
Rheum rhaponticum, *Phytophthora parasitica* susceptibility, 290
Rhizoctonia, aerial blight of Lespedeza, soil source of inoculum, 293
ethyl mercury phosphate, control by in gladiolus corms, 154
foliage blight, cucumber, 290
Lysol and ferbam, control by, in gladiolus corms, 5
solani, bean seeds, 483
conifer damping-off control, 466
cotton dusting for control, 292
flax seedling blight, *Corticium praticola* as perfect stage, 488
gladiolus corm lesions, 149
inhibition by soil organisms, 586
lettuce, control, 475
methyl bromide control, 376
on potato in India, 645
potato stems, control, 475
tobacco seedlings, damping-off control, 125
tomato control, 475
turf brown patch, factors influencing, 484
sp., sugar beet seedlings black rot, 300
spp., celery, control in seedbeds, 293
web-blight symptoms on beans, 290
Rhizoglyphus rhizophagus, gladiolus corms, 150
Rhizopus nigricans, fungicide control on peach, 407
on soybean, 292
survival on cotton seeds in storage, 220
spp., on peanuts in market, 474
tritici, effect on respiration of sweet potato root, 222
Rhopalosiphum spp., see also Insects, aphids, cereal yellow-dwarf virus vectors, 132, 309, 542
Rhubarb, ringspot-like virus, 590
RICE, ROBERT V., 482
RICH, A. E., (109)
RICH, SAUL, (476), 482
RICHARDS, M. C., 109
RIEMAN, G. H., (490)
RIKER, A. J., (92), (441), (448), (475), (487)
Rimocidin, see Antibiotics
RODENHISER, H. A., 558
Root rot on pine seedlings, control by soil fumigation, 81

Rose bengal, in isolation of Fusarium spp. from plant tissues, 700
Rose, black spot, fungicide-insecticide mixture for, 406
rosette virus, 218
Ross, A. Frank, 1
Roth, Elmer R., (475)
Rotylenchulus reniformis, see Nematodes
Rowell, J. G., 654
Rubus spp., apple flyspeck, 586
Rust, bean, pressure effects on, 70
flax, epidemiology in North Central States, 624
physiologic race survey from 1931, 625
races, genetic characters, 626
varieties resistant to, fungus races interaction, 625
fungi, pressure effects on, 70
germ tube development, zinc and silicon or gelatin requirement, 483
mint, method of evaluating severity and control, 587
oak, pycniospore stage on pines, 488
oat crown, effect of temperature and age of plant on varietal reaction to, 484
on gall tissue of *Juniperus virginiana* in culture, 360
peppermint, oil gland destruction and leaf drop, 587
physiology, 360
snapdragon, pressure effects on, 70
sugar beet resistant varieties, 300
sunflower, pressure effects on, 70
wheat black stem, races of pathogen manner of infection of resistant varieties, 485
breeding for resistance, "composite variety," 467
leaf, distribution in Oklahoma, 294
leaf and stem, control by, and phytotoxicity of chemotherapeutants, 496
races, 294
stem, breeding for resistance, 472
infection type, Kubanka, 481
races, pathogenicity on Kenya varieties, 474
races, resistance, 472
seedling and adult changes in resistance, 468
spore shower, origin and extent, 471
white pine blister, distribution, effect of microclimate, 487
yellow, races of, on wheat crosses, 471
Rye, *Helminthosporium* spp., reaction to isolates of, 517
yellow-dwarf virus, 131

Saccharine, wheat stem rust, 661
Saintpaulia ionantha, root-knot nematode, 406
Sakimura, K., 217
Salicylic aldehyde, effect on virulence of Pythium root rot pathogen, 289
Sallans, B. J., (701)
Sambucus glauca, apple flyspeck, 586
Sasser, J. N., 483
Scab, cucumber, breeding for resistance to, 215, 257
Scheffer, R. P., 116, (472), 483, (535)
Schexnayder, C. A., (289)
Schlegel, David E., 89, 206, 589
Schnathorst, William C., 483
Schneider, Henry, (51)
Schroeder, H. W., 401
Schuster, M. L., 483
Sclerospora sp., on sugarcane, 289
Sclerotinia fructicola, effect of fungicides on spore germination and hyphal growth, 476
polyphenol oxidase relation to fungitoxicity, 483
S. laxa, pathogenicity compared on various hosts, 589
gladoli, corm lesions, 148
laxa, 589
sclerotiorum, bean, effect of storage of sclerotia on viability, 520
bean infection by perfect stage, 520
bean, occurrence and source of infection, 519
soil organisms on, 471
spp., fungitoxicity of sodium and potassium salts of 2-benzothiazolythioglycolate to, 43
Sclerotinia fructicola—*continued*
methyl bromide control, 376
Sclerotium bataticola, effect of temperature and moisture on growth, 634
linear growth rates, 633
delphinii, soil organisms on, 471
rolfsii, calcium inhibition of, 363
disintegration in alfalfa meal-soil mixture, 366
nitrite, toxic effect, 366
pathogenicity of isolates on peanuts, 289
soil microflora antagonism toward, 366
sweet potato circular spot, 432
spp., tobacco seed-bed incidence, 125
Scolytidae, see Insects
Scott, C. Emlen, (218)
Seed treatment, see also under Fungicides
barley loose smut control, 313
grasses and legumes, 487
mercury fungicides, vapor action of, 465
oat, 290
onion smut, thiram, 596
peanut, 291
pelleting, conifer damping-off, 466
tomato, damping-off, 291
wheat loose smut, 315
Segall, R. H., 483
Senseney, C. A., (687)
Septoria, barley leaf blotch, 473
apii-graveolentis, celery resistance to, 254
linicola, dissemination, 468
tritici, 294
Serological test for tobacco mosaic virus, 77
rabbit antisera, TMV strains, antigenic relationships, 589
with *Pseudomonas* spp., 412
Shallots, pink root control, 294
Shands, R. G., (472)
Sharp, E. L., 483
Shaw, Charles Gardner, (181)
Shaw, R. H., (317)
Shay, J. R., (156), 483
Sherbakoff, C. D., 395
Shurtleff, M. C., 110, 484
Siang, W. N., (219)
Siegel, Albert, 589, (589)
Silberschmidt, Karl, 304
Sill, W. H., Jr., 484
Silverborg, S. B., 20, 699
Simmonds, P. M., (478), 484, (701)
Simons, John N., 29
Simons, M. D., 484
Sitanion hystrix, 310
Skiles, R. L., 409
Skotland, C. B., 484
Slykhuis, John T., 484, 537
Smith, D. H., (474)
Smith, Floyd F., (404), 406
Smith, F. G., (483)
Smith, M. A., 583
Smith, W. L., Jr., (50), 406
Smut, barley, 393
covered, inoculation method, 697
ergot associated with, 461
fungi, nomenclature, system of classification for, 181
grass head, differentiation of races, 200
striped, host range and distribution, 527
specialized varieties and physiologic races of pathogen, 528
symptoms and pathological histology, 527
oats, control, 393
covered, inoculation method, 697
onion, control, thiram seed treatment, 596
wheat, control, 315, 461
loose, distribution in Oklahoma, 294
races of pathogen, 681

Snapdragon, cucumber mosaic virus strains on, 336
rust, pressure effects on, 70
Snow-mold, see Alfalfa, crown rot
SNYDER, WILLIAM C., (470), (586)
Sodium salts of dimethyl dithiocarbamic acid and of 2-mercaptobenzothiazole, see Fungicides, Vancide 51
Sodium salt of dinitro ortho cresol, see Fungicides, Elgetol
Soil, antibiotic production in, 108
Soil composition, injury from methyl bromide effected by, 489
Soil fumigants, general, citrus nematode, 466
fungi-nematode complex in celery seedbeds, 293
orange, effect on growth and yield, 465
pine root rot, 81
strawberry red stele control tests, 590
Verticillium control and methods of evaluating effects, 593
Soil fumigants,
allyl bromide, 81
CBP-55 (chlorobromopropene), 81, 293, 593
root rot, pea and bean, 590
chloropicrin, 81, 593
conifer seedling disease complex, root-rot and damping-off, 469
tobacco, effect on growth and *Pythium ultimum* infection, 590
DD (dichloropropene-dichloropropane), 81, 293, 466
nematode detection, 260
sting nematode control, 291
sweet potato circular spot, 433
dichlorobutene, 81, 293
Dowfume W-40, sting nematode, 438
EDB (ethylene dibromide), 81, 293, 466
effect on nematode populations, 474
Fusarium wilt-nematode complex, cotton, 293
nematode detection, 260
pine seedling root rot control, 81, 474
sting nematode control, 291
sweet potato circular spot, 433
formaldehyde, root rot control, pea and bean, 590
Larvacide, see chloropicrin
methyl bromide, 81, 293
effect on various plant species, 376
injury, carnation and chrysanthemum, 489
injury, effect of soil composition and other treatments upon, 489
various fungi, 376
Soil fumigation, apparatus for, 81, 590
Soil microflora, antagonism toward *Sclerotium rolfsii*, 366
associated with *Verticillium* sclerotia, 594
effect of carbon and nitrogen on, 365
effect of organic matter on, 367
Fusarium vasinfectum growth in presence of, 482
non-pathogenic organisms for reinfestation of sterilized soil, 586
on sclerotia, 471
Soil moisture, effect on growth and survival of *Fusarium oxysporum* f. *cubense*, 499
Soil temperature, effect of air temperatures and sky condition on, 573
effect on cabbage virus A in *Nicotiana* spp., 550
effect on heat canker of peanuts, 573
effect on turnip virus 1 strains in *Nicotiana* spp., 482
related to Botrytis neck rot of gladiolus, 167
root and stem, internal temperatures compared, 317
Soil treatments, conifer damping-off, 466
effect on growth of tobacco and strawberry, 590
soybean meal in wheat root rot control, 701
Soil type, *Fusarium oxysporum* f. *lini*, effect on virulence, 493
relation to moisture in growth and survival of *Fusarium oxysporum* f. *cubense*, 500
Soja max, see Soybean
Solanum demissum, late blight, races of pathogen on, 480
Phytophthora infestans races in Mexico, 454
melongena, see Eggplant
Solanum—*continued*
nodiflorum, potato virus Y on, 217
spp., *Heterodera* sp. on, 106
Phytophthora infestans on, 456
parasitica, resistance to infection by, 290
Synchytrium endobioticum on, in Mexico, 480
tobacco-etch virus on, 568
virus, resistance, 245
Sorghum spp., 310
Sorosporium spp., nomenclature, changes in, 183
SOWELL, GROVER, JR., 485
Soybean, *Alternaria* spp. on, 293
Ascochyta spp. on, 469
bacterial blight, inoculum dilution and host relation, 468
persistence of pathogen, 191
bacterial pustule, field and laboratory tests on persistence of pathogen, 189
wildfire, effect of presence of, 191
Cercospora kikuchii on, 292
purple stain, seed, 477
downy mildew race 4 of pathogen, 292, 460
fungi associated with seeds and pods, 292
necrotic-lesion strain of yellow bean mosaic virus on, 46
root-knot nematode, clean fallow for control, 405
seed treatment, autumn, 476
sting nematode, 295, 435
wildfire, bacterial pustule, effect of presence of, 191
field and laboratory tests on persistence of pathogen, 189
SPANGELO, L. P. S., 345
Sphacelotheca spp., nomenclature, changes in, 183
SPILKER, OREN W., 485
Spinach, clover-root nematode on, 606
cucumber mosaic virus strains on, 334
downy mildew, pressure effects on, 70
effect of air temperature and nitrogen supply on injury by olefinic peroxide fumigants, 588
tomato aspermy virus on, 404
Spinacia oleracea, see Spinach
Sporisorium spp., nomenclature, changes in, 183
Sporotricum sp., in bean seeds, 483
Spot anthracnose, Bouchea in Cuba, 406
SPRAGUE, V. G., (642)
SPROSTON, THOMAS, JR., 110
Sprout inhibitors, Fusarium dry rot of potatoes, 95
Squash, disease resistant hybrids, 257
summer, Phytophthora blight, 469
SR-406, captan, coined name, 58
STAGE-SMITH, R., 589
Stagonospora maculata, effect of periods of high and low humidities on development, orchardgrass, 642
STAHMANN, MARK A., (473), (482), (535)
STAKMAN, E. C., (470), (471), (474)
STALEY, JOHN M., (341)
STARR, C. H., 439
STEERE, RUSSELL L., 485
Stemphylium botryosum, alfalfa leaf spot, 477
conidial stage of *Pseudoplea briosiana* on alfalfa, 584
onion brown blotch, 411
loti sp. nov., birdsfoot trefoil, symptoms, 577
cardinal temperatures, pathogenicity and overwintering, 579
host range and cross-inoculation studies, 579
sarcinaeforme, polyphenol oxidase relation to fungitoxicity, 482
sp. fungitoxicity of sodium and potassium salts of 2-benzothiazolylthioglycolate to, 43
in bean seeds, 483
STESSEL, G. J., 23, (61), 65
STEVENSON, F. J., 245
STEWART, R. N., (406)
Stizolbium deeringianum, see Bean, velvet
Stone fruit, green ring mottle virus, 326, 329
latent virus complex, cherry flower inoculum, 479

Stone fruit—*continued*
necrotic ringspot virus, 325, 329
viruses in cucurbit hosts, effect of extracts on infectivity, 328
infectivity and longevity in cucurbit hosts, 324
quantitative method of assay on cucumber, 175
STONER, WARREN N., 293, 611
STOVER, R. H., 499, 700
Strawberry, gray mold control, 482
leaf blight control, 471
leaf scorch, photographic infection scales for roguing susceptible seedlings, 345
leaf spot, photographic infection scales for roguing susceptible seedlings, 345
red stele, soil fumigation and fungicide tests, for control, 590
sting nematode, 295, 435
STREETS, R. B., 589
Strepogenin, lycomarasmin inhibition of, 233
Streptomyces lavendulae, 359
scabies, 355
potato resistance to, 250
sp., antimycin A (A-102) produced by new isolate of, 478
spp., amino acid effect on melanin in, 357
melanin production and growth in, 355
toxin formation, 356
venezuelae, synthesis of chloramphenicol by, 478
Streptomycin, see Antibiotics
STROUBE, WILLIAM H., 293
STRUBLE, F. BEN, (290)
Sugar beet, breeding for disease resistance, 297, 301
breeding techniques, 301
black root, causal organisms and resistance, 300
crop: size, geographical distribution and economic importance of, 297
curly-top virus strain, nature of injury, 588
on potato, synonym green dwarf, 684
resistance to, 298, 588
virus yellows relationship, 587
downy mildew, non-bolting resistant varieties, 300
leaf spot, resistant varieties and hybrids from inbreds, 299
seed production enterprise, 298
varieties, combined disease resistance, 300
virus yellows, 301
virus yellows, curly-top relationship, 587
varieties, resistant, 302, 405
Sugarcane, growth-retarding disease in Louisiana, 289
nematodes, associated with diseased roots of, 289
Pythium root rot, salicylic aldehyde, effect on virulence of pathogen, 289
Sclerospora sp. on, 289
Sugars, carnation bacterial wilt studies, 587
Sulfa compounds, wheat rust, control and phytotoxicity, 497, 661
Sunflower, rust, pressure effects on, 70
tumor tissue *in vitro*, culture of powdery mildew on, 343
Sun-hemp, charcoal rot, pycnidial stage of fungus on, 608
SWAEBLY, MARY ANN, 485
SWANK, GEORGE, JR., 285, (293)
Sweet pea, cucumber mosaic virus strains on, 336
Sweet potato, black rot, differences in susceptibility of varieties to, 78
internal cork disease, varietal reaction to, 290
nematode, root-knot, resistance to, 290
Sclerotium rolfsii, circular spot, 432
sting nematode, 435
storage temperature, relation to internal cork development, 50
Sweet william, carnation Fusarium wilt on, 465
SYLVESTER, EDWARD S., (29), 209, 541
Symphoricarpos albus, *Phytophthora parasitica*, susceptibility to infection by, 290
Verticillium albo-atrum on, 588
Synchytrium endobioticum, potato resistance to, 251
wild potatoes in Mexico, 480
Syngonium auritum, black cane rot, 586
SZKOLNIK, MICHAEL, (109)

Tagetes signata, bayberry-yellows virus, 18
TAKESHITA, R. M., 485
TAMMEN, JAMES, 485
Tangerine, wood rot and canker associated with *Fomes applanatus*, 99
Taphrina coerulescens, oak, control, 472
TAPKE, V. F., 162, 407
TARJAN, A. C., 485, 486
Taxonomy, see also Nomenclature, *Erwinia chrysanthemi*, 525
Pseudoplea briosiana, 584
Stemphylium botryosum, 584
TAYLOR, GORDON S., 486
TAYLOR, JACK, 268
Techniques, alfalfa resistance to leaf spot, rapid method of measurement, classification of plants, 651
apple seedlings in greenhouse, testing of fungicides, 109
autoradiography of histological sections with labeled toxins, 237
bacteriophage in differentiating types of *Xanthomonas pruni*, 27
barley loose smut viability prolonged by storage temperatures, 407
chromatographic analysis of extracts of plants infected with radioactive pathogens, 237
chromatographic separation of 1-4 naphthoquinones, 110
culturing obligate parasites on excised tissue *in vitro*, 343
equipment, fungicide-insecticide dusting, seedling diseases and insect pests, 480
soil fumigation, 590
evaluating effects of soil fumigants on *Verticillium*, 593
field inoculation testing tobacco for black shank resistance, 289
freezing, causing breaking of TMV particles, 482
greenhouse test for susceptibility of pear seedlings to fireblight, 158
heat-ultra method for purification of TMV for spectrophotometric assay, 89
identifying potato seedlings immune from virus X, 487
inoculation, barley and oats seed with smut spores, 697
Fusarium oxysporum f. *lini*, 491
pear seedlings with *Erwinia amylovora*, 156
inoculum dilution in testing for resistance, soybean to bacterial blight, 468
flower and leaf compared, stonefruit viruses, 480
plant disease fungi, large quantity preparation, 401
insect injection, 388
isolation of fungi from woody tissue, instruments, 403
medium for tissue culture, fungi, 360
nematocides, assay, 264
nematodes, detection in the field, 259
diagnosis by analysis of mineral elements of plant parts, 260
laboratory culture for teaching and research, 105
laboratory preservation, 262
soil sampling for detection, 261
staining plant tissues for detection, 261
temporary and permanent mounts, preparation of, 262
paint spray gun, mass inoculation with potato virus X, 487
partial vacuum method of inoculating seedlings, *Ustilago zeae*, 654
pathogenicity of biotypes of *Ustilago zeae*, method to determine, 654
preservation of cultures under oil, 407
proving pathogenicity of vivotoxins, 229
pure culture maintenance and inoculation procedure, *Pseudopeziza medicaginis*, 651
radio carbon C^{14} in medium for oak wilt fungus, 490
radioisotopes, selection for toxin assay, 236

Techniques—*continued*
radio phosphorus in Czapek's medium, 478
rose bengal and streptomycin in isolation of *Fusarium* spp., 700
rust, mint, evaluating severity and control, 587
scale model to estimate reduced water flow in Fusarium wilt, 488
serological tests, rabbit anisera, TMV strains antigenic relationships, 489
sheep blood for tobacco mosaic virus, 77
spores in soil, population estimation, *Helminthosporium sativum*, 478
spraying, mist and hydraulic compared, 465
staining vascular elements, 117
strawberry, photographic selection scales, roguing seedlings susceptible to leaf spot and leaf scorch, 345
sugar beet breeding for resistance, 301
temperature of diseased and healthy leaves measurement, 676
tests for fungicidal solution strength for bulb dips, 598
toxins demonstrated in tomato Fusarium wilt by intergeneric grafts, 470
ultra violet absorption in determining concentration of purified TMV, 488
virulence retention in lyophylized culture of *Pseudomonas caryophylli*, 406
virus assay, quantitative method for stone-fruit viruses on cucurbits, 175
virus cherry seed transmission, quick method, 467
virus concentration estimation, biological assay and ultraviolet absorption method compared, 589
virus inoculum, flower tissue, 479
virus purification, electrophoresis and density gradient centrifugation, 467
wheat breeding for rust resistance, "composite variety," 467
woody tissue, instruments for isolation of fungi, 403
Temperature, air and hot water, lethal to *Verticillium albo-atrum*, 589
air and soil, effect on turnip virus 1 strains in *Nicotiana* spp., 482, 550
air, effect on injury of olefinic peroxide fumigants on herbaceous plants, 588
bayberry-yellows virus control, 18
effect on, bacteriophage plaque formation, 28
cabbage yellows and Fusarium wilt of radish, 277
development of Fusarium wilt on flax, 492
disease development in gladiolus corms, 141
Erysiphe graminis tritici on seedlings in greenhouse, 290
gall development, corn smut, seedlings, 656
growth and disease ratings, pear seedlings in greenhouse, 160
hydrogen cyanide production by alfalfa crown rot pathogen, 581
lesion formation by potato virus Y on *Physalis floridana*, 2
longevity of spores of oak wilt pathogen, 476
longevity of *Ustilago nuda*, 407
oat varietal reaction to crown rust, 484
production and survival of sporangia of *Phytophthora infestans*, 505
radial growth of *Ustilago striiformis* varieties, 534
reaction of Kubanka wheat to races of *Puccinia graminis tritici*, 481
survival of *Colletotrichum gossypii* on cotton seeds, 220
susceptibility of barley to powdery mildew, 162
susceptibility of oat hybrids to stem rust, race 7, 472
wound periderm and cuticle, gladiolus corms, 141
heat-ultra method of purification of TMV for spectrophotometric assay, 89
hot water and air, lethal to *Verticillium albo-atrum*, 589
of diseased and healthy leaves, relation to respiration, 679
of diseased, injured and healthy leaves compared, 675

Temperature—*continued*
of host tissue as affected by parasites, 221
of *Pythium ultimum*, internal plant, related to damping-off by, 318
of root and stem, internal, compared with soil, 317
relation to internal cork development, sweet potato roots, 50
soil and air, effect on turnip virus 1 strains in *Nicotiana* spp., 482, 550
soil, effect on gladiolus reaction to Botrytis neck rot, 167
—time gradients control, mushroom-spawn nematode, 404
Terramycin, see Antibiotics
Tetrachloronitrobenzene, effect of, on Fusarium dry rot in potatoes, 95
Tetrachloro-para-benzoquinone, see Fungicides, chloranil
Tetramethylthiuram disulfide, see Fungicides, thiram
Thecaphora spp., nomenclature, changes in, 183
Thelephoraceae, wood rotting by, 21
Thielaviopsis basicola, tobacco breeding for resistance to, 239
Thiolutin, see Antibiotics
Thiouracil, tobacco mosaic virus, inhibition of, related to light, 555
THIRUMALACHAR, M. J., 527, 608, 645
THOMAS, H. EARL, 218
THOMAS, R. REX, 11, (407)
THOMAS, W. D., JR., (587)
THOMPSON, G. E., 293
THORNBERRY, H. H., 486
THURSTON, H. DAVID, 486
THURSTON, H. W., JR., (108)
TIFFANY, LOIS H., 486
Tilletia asperifolia, spore germination of, 219
brevifaciens, spore survival, exposed on soil, 487
caries, conditions affecting resistance of wheat, 473
differential survival, *T. foetida* in mixture on host, 558
hybridization, *T. foetida*, 559
pathogenic biotypes, 399
sex compatibility factors, 322
sporidial fusion in culture, 322
foetida, differential survival, *T. caries* in mixture on host, 558
hybridization, *T. caries*, 559
spp., nomenclature, changes in, 183
taxonomy, spore germination related to, 219
TIMIAN, ROLAND G., 487
TIMS, E. C., 294
TINLINE, R. D., 487
Tobacco, bacterial wilt, resistance to, 240
black root rot, resistance to, 239
black shank, field inoculation effectiveness for testing resistance, 289
root-knot nematode effect on resistance, 483
tomato fruit rot incited by, 290
toxin production, 294
blue mold control, 486, 616
distribution, geographical, 616
oospores of pathogen, germination, 701
overwintering of pathogen, 616
spore showers, 616
zoospores, infection by, 701
breeding for resistance, 239, 243, 469
copper sulphate toxicity symptoms on seedlings, 125
cyst nematode, host preference, 106
damping-off of seedlings, control of, 125
disease resistance, 239, 243, 469
etch virus, 564
Fusarium spp., 590
mosaic virus assay, see also under Viruses, 77, 89, 484, 485, 589
effect of organic compounds on multiplication, 589
effect on temperature of leaves, 679
growth curve of host in relation to concentration curve of virus, 488

Tobacco—*continued*
host nutrition in relation to virus multiplication, 488
pressure effects on, 70
radiophosphorus inhibition of, 206
resistance to, 242, 475
strains, 587
thiouracil, control and plant injury, 555
necrosis virus, 25, 590, 679
Pythium ultimum on roots, 590
radiophosphorus injury, 207
resistance to combination of diseases, 240
ring spot virus, 70, 485, 687
soil treatment effect on growth, 590
thiouracil up-take by leaves, related to light, mosaic virus studies, 556
Tomato, awl nematode, 296
bacterial canker, seed transmission, and control, 473
buckeye rot, susceptibility to, 290
bushy-stunt virus strains of, 485
damping-off seed treatment, 291
disease resistance, effect of growth-regulators, 137
early blight, control, 25, 478
Fusarium wilt, chemotherapy, 43, 137
copper ion effect on, 283
enzymes role in etiology of, 470
ethyl alcohol relation to, 663
ethylene, epinasty, 470, 663
ethylene production in infected plants, 664
fungus distribution, 116
iron ion effect on, 283
lycomarasmin inhibitors effect on, 283
lycomarasmin role in, 284
mechanism of vascular discoloration, and wilting, 470
metabolites production of vascular browning, 490
nutritive sprays effect on, 466
osmotic properties of leaf parenchyma in diseased plants, 619
pectic enzymes, 472, 535
physiology, 116
8-quinolinol effect on, 283
reduction of water flow estimated by scale model, 488
resistance affected by metabolic inhibitors, 483
toxin demonstrated by intergeneric grafts, 470
transpiration, velocity, 621
vascular reactions to, 116
golden nematode, cytological effects of, 471
internal temperatures of root and stem compared with soil, 317
iron deficiency, effect on response to lycomarasmin, 116
late blight, resistant varieties reaction to virulent race of pathogen, 344
sporangia production and survival, 505
leaf mold, varieties immune from, 109
lycomarasmin effect on, 195, 281
nutrition, effect on susceptibility to *Alternaria solani*, 476
root-knot control, 465, 485
root-knot, resistance to, 290
root rot control, 465
stubby-root nematode, 296
tobacco cyst nematode, 106
toxic metabolites, effect on, 116
Verticillium, soil fumigation tests and control, 595
virus, apparently new, 480
Toxic metabolites, lycomarasmin, 116
vascular browning production, 490
Toximycin, see Antibiotics
Toxins, abolition of regulatory mechanisms in hosts, 221
defense reaction and pathogen, chemical aspects, 353
2,4, dinitrophenol, effect on respiration compared, 224
effect on phosphorus metabolism, 225
modes of action, 355
nature of defense reaction to, 351
physiology of formation in microorganisms, 355
produced by *Fusarium*, 352
Toxins—*continued*
production by host—parasite interaction in tomato Fusarium wilt, 470
production in relation to tobacco black shank, 294
proteins, carnation wilt studies, 587
radioisotopes, use in detecting toxin and host tissue reactions, 236
respiration, effect on, 221
tobacco black shank, adsorption on charcoal, 294
tomato Fusarium wilt, effect upon species grafted to tomato stocks, 470
vivotoxins, 229
wound reactions to, 352
Toxoptera graminum, see Insects aphids, greenbug
Transpiration, magnesium and potassium salts effect on tomatoes in relation to Fusarium wilt, 466
N-trichloromethylthio tetrahydrophthalimide, see also Fungicides, captan, coined name, 58
Trichoderma spp., corn seedling blight, associated with, 477
in soil, inhibitory to damping-off of peppers, 586
Trichodorus sp., see Nematode, stubby-root
Trichothecium roseum on peanuts in market, 474
2,3,5 triiodobenzoic acid, Fusarium wilt of tomato, effect upon, 137
metabolism of host, effect upon, 139
Triticum dicoccum, wheat stem rust, breeding for resistance, 472
timopheevi, powdery mildew resistance to, 290
Tritium, see radioisotopes
Tropaeolum majus, see Nasturtium
TROXEL, A. W., (477), 590
TRUE, R. P., 487
TRYON, E. H., (487)
Tuburcinia spp., nomenclature, changes in, 183
Tumors, see Galls
Turf, see also Grasses, brown patch (*Rhizoctonia solani*) factors influencing, 484
brown patch, susceptibility of varieties to, 110
Curvularia "fading-out," 109
TURNER, W. F., (691)
Turnip, virus 1 strains, effect of soil and air temperatures on in *Nicotiana* spp., 482, 550
Tylenchorhynchus, see Nematodes
TYLER, L. J., 487
Tyloses formation in oak wilt, 451
TYNER, L. E., 313

Ultra violet, absorption method for estimation of TMV concentration, 589
spectra, tobacco mosaic virus, groups of strains compared, 587
Uredo spp., nomenclature, changes in, 183
Urocystis spp., nomenclature, changes in, 183
Uromyces betae, sugar beet resistant varieties, 300
fallens, effect on respiration of clover leaves, 223
phaseoli, pressure effects on, 70
Usnic acid, see Antibiotics
Ustilagidium spp., nomenclature, changes in, 183
Ustilago bullata, differential grasses, 201
races, new, 200
x *U. striiformis hordei*, inheritance of spore and sorus characters, 547
nuda, control on barley and wheat, seed treatments, 313
differential barley varieties, 407
longevity, 407
races, 369, 407
races, spore germination characteristics, 482
temperature effect on longevity, 407
spp., ergot associated with, 461
inoculation method, barley and oats, 697
nomenclature, changes in, 183
striiformis, effect of temperature on radial growth, varieties, 534
horedi, x *U. bullata*, inheritance of sorus and spore

Ustilago striiformis—*continued*
characters, 547
host range, 527
nomenclature, 530
varieties and races, 528
varieties, spore germination, cultural characters and cytology compared, 527
tritici, effect on respiration of wheat leaves and stems, 223
races, 681
races, spore germination characteristics, 482
wheat germ plasm effect on prevalence of races, 683
zeae, amino acids synthesized by, 470
partial vacuum method of inoculating corn seedlings, 654
pathogenicity of biotypes, method to determine, 654
spore germination, effect of potassium phosphate, 478

Vaccinium australi, see Blueberry
VALLEAU, W. D., 616
VALLEGA, JOSE, (471), 487
VAN ARSDEL, E. P., 487
VANTERPOOL, T. C., 488
VATTER, A. E., (486)
VAUGHAN, EDWARD K., (589), 590
Vaughan, Richard English, Biographical sketch, 349
Venturia inaequalis, amino acids as nitrogen sources, 481
races 2 and 3, physiologic specialization, 483
Verticillium albo-atrum, control, 376, 475, 594
eggplant, isolates, 466
Fusarium associated with, on potatoes, 108
Gossypium spp., reaction to, 489
host range, 588
internal brown spot associated with, on potatoes, 108
microflora and nematodes associated with sclerotia in soil, 594
pathogenicity of isolates on peppermint, 588
pepper, isolates, 466
"pink eye" associated with, on potatoes, 108
Pseudomonas fluorescens, associated with, 108
soil fumigation tests, methods of evaluating effects, 593
storage behavior of potatoes infected with, 108
temperatures lethal to, hot water and air, 589
Vicia americana, see Common vetch
faba, see Bean, broad
Vigna spp., see Cowpea
Vinca rosea, see Periwinkle
Viruses, general, acquisition threshold periods and vector efficiency in 4 species of aphids, 30
concentration in plant tissue, biological assay, ultraviolet absorption method compared, 589
effect on temperature of infected leaves, 679
electron microscopy of virus particles, 486
infectivity inhibited by synthetic polypeptides, 473
inoculum, flower tissue for isolations, 479
loss of insect vector transmissibility by, 466
mosaic virus complex on Ladino clover, 271
phony disease, peach and plum, transmission of virus from tops of peach and plum trees, 691
purification, electrophoresis and density gradient centrifugation, 467
quantitative method of assay of stone-fruit viruses on cucumber, 175
Viruses,
alfalfa mosaic, differentiation of strains, 479
local lesion strain, 275
strains, host range, 271
systemic strain, 275
thermal inactivation, 274
tomato varieties, 479
Aureogenus spp., insect vectors, 9
loss of insect vector transmissibility, 466
purification, 467
bayberry-yellows, 15
bean 2, gladiolus, 111

Viruses—*continued*
pea, resistant variety, New Era, 473
bean southern mosaic, serological test for, 77
bean yellow mosaic, age of plant and method of inoculation, 473
mild and necrotic strains, 275
necrotic-lesion strain, cross-protection, 45
on gladiolus, 114
on tobacco and bean varieties, 11
red clover symptoms, 470
severe-yellow-mosaic strain, 11
strains, host range, 271
Brassica nigra, age, effect on transmission of, 212
age of plant, effect on plant susceptibility, 211
aphid-host plants interaction variation, 209
aphid transmission, 542
darkness, effect on plant susceptibility, 210
host range, 541
on *B. juncea*, 209
properties: thermal inactivation, dilution end-point, longevity *in vitro*, 541
site of inoculation, effect of, on transmission by green peach aphid, 209
source, effect on transmission of, 211
cabbage A, effect of air and soil temperatures on, in *Nicotiana* spp., 550
cabbage black ring, effect of air temperature on, in *Nicotiana* spp., 550
cabbage mosaic, Badger Market, new variety, resistance to, 650
camellia, transmitted, 293
cauliflower 1 group, properties, 545
celery western mosaic, aphid transmission, 29
cereal yellow-dwarf, aphid vectors, 132, 309, 312
control, 312
epiphytotic, conditions favoring, 309, 312
grass species immune from, 310
host range, 135, 309
red-spot mosaic, compared, 135
symptomless carriers, 309
cherry complex, flower tissue inoculum, 479
cherry green ring mottle, 481
cherry necrotic ring spot, quick test for seed transmission, 467
cherry recurrent chlorotic ring pattern, 325
cherry ring spot, effect on growth of nursery trees, 480
cherry tatter leaf, infectivity and longevity, of inoculum, 326
cherry yellows, effect of aging on infectivity of inoculum from vegetable marrow and pumpkin, 325
infectivity and longevity in cucurbit hosts, 324
inhibitors of infectivity of isolates of, 330
quick test for seed transmission, 467
transmission from peach understock to Japanese plum, 108
Chlorogenus myricae, 19
chrysanthemum stunt, *Marmor chrysanthemi*, proposed name, 406
properties: dilution end-point, longevity *in vitro*, thermal inactivation, 406
citrus tristeza, 51
crucifer, aphid-transmissible, radish 1 group, 545
thermal inactivation and dilution end-point of strains, 545
cucumber mosaic, gladiolus, 111
inoculum, gladiolus flower tissue, 479
on Japanese barberry, 489
strains from alfalfa and pepper on pepper, 333
grape degeneration, related to Pierce's disease, 611
yellow-headed sharpshooter vector, 612
grape Pierce's disease, leafhopper vector, 293
grape yellow mosaic, 475
horseradish mosaic, effect of air temperature on, in *Nicotiana glutinosa*, 550
Lethum australiense, purification, 467

Viruses—*continued*

Myrica virus 1, 19
nasturtium mosaic, Brazilian compared with other nasturtium mosaics, 307
distribution of, in inoculated plants, 304
Lycopersicum virus 3 relation to, 306
properties, 307
strains of, 307
symptoms on naturally and experimentally infected plants, 304
oat Siberian mosaic and wheat striate mosaic compared, 539
panax ringspot, symptoms and transmission, 643
pea Wisconsin streak, pea aphid vector, 484
peach mosaic, spread related to age of infection, 585
peach necrotic ringspot, 108
peach western X, ninhydrin-reacting substance in leaves, 586
physiological changes induced by, 481
respiration of healthy and infected foliage, 481
peach yellow leaf roll, leafhopper transmission studies, 562
potato A, strains, 479
Myzus persicae vector, 479
potato curly-top, strain, 587
synonym green dwarf, 684
potato leaf roll, 248
on tomato, effect upon response to naphthaleneacetic acid, 109
potato mosaic, 247
potato net necrosis, 249
potato X, method for identifying immune seedlings, 487
potato Y, aphid transmission, 217
mechanical inoculation vs. aphid inoculation, 469
Physalis floridana, 1
properties and host range, Hawaii, 217
prune dwarf, infectivity and longevity, of inoculum, 326
radish 1 group, 545
red clover vein-mosaic, age of plant and method of inoculation, 473
incidence and host range, 473
pea stunt, 473
rhubarb ringspot-like, 590
rose rosette, 218
spotted wilt on tomato in Hawaii, 217
stone-fruit, in cucurbit hosts, effect of extracts on infectivity, 328
infectivity and longevity in cucurbit hosts, 324
stone-fruit, green ring mottle, 326
stone-fruit latent complex, by cherry flower inoculum, 479
stone-fruit necrotic ring-spot, 325
sugar beet curly-top, strain, nature of injury, 588
yellows relationship, 587
sugar beet yellows, 301
curly-top relationship, 587
tobacco-etch and TMV compared on pepper species, 564
Chenopodium album, new local lesion host, 568
tobacco mosaic, bean acquired resistance, 490
breaking of particles caused by freezing, 482
classification of strains, 589
effect of organic compounds on multiplication, 589
electron microscopy *in vivo*, 484
electrophoretic mobility, groups of strains compared, 587
growth curve of host in relation to concentration curve of virus, 488
host nutrition in relation to virus multiplication, 488
in tissue culture of tobacco, synthetic media, 475
isoelectric point, groups of strains compared, 587
mutants, serological tests for, 77
nitrogen, nutrition effect on concentration of, in tomato, 477
nutrition of host tissue in virus activity, 575
particles associated with granules in plant tissue revealed by electron microscopy, 484
pressure effects on, 70
purification of, by heat-ultra method for spectrophotometric assay, 89
radiophosphorus effect on, 206
rosette strain, serological test for, 77
serological test for, 77
serological tests for antigenic relationships, strains, 589
size of particles, 482
strains, continous variation, 587
strains, effect of temperature on toxicity, 587
thiouracil control and plant injury, 556
thiouracil inhibition of, related to light, 555
ultra-violet absorption spectra, groups of strains compared, 587
tobacco necrosis virus, effect on temperature of bean leaves, 679
strain, host adaptation, bean and petunia, 590
tobacco ring spot, activity and particle sedimentation and counts from electron micrographs, 688
electron microscope studies, 687
electrophoretic fractionation and mobility, 687
hexagonal particles, concentration and purification, 485
on gladiolus, identification of strains, 112
particle size and shape, 688
pressure effects on, 70
purification, 485, 687
tomato, apparently new, described, TMV compared, 480
tomato aspermy, aphid vectors, on chrysanthemum and tobacco, 404
on various plant species, 404
tomato bushy-stunt strains, 485
tomato ringspot on *Hydrangea macrophylla*, 404
turnip 1, effects of air and soil temperature on, in *Nicotiana* spp., 550
host nutrition and virus multiplication in *Nicotiana* spp., 669
properties, 545
strains, effect of soil and air temperatures on, in *Nicotiana* spp., 482
variegation, leaf and flower, camellia, graft transmitted, 293
wheat Russian mosaic and striate mosaic compared, 539
wheat streak mosaic, *Aceria tulipae* vector, 484
effect of wind and blowing soil on infection, 484
elongated particles associated with, electron microscopy, 458
wheat striate mosaic, host range, 539
leafhopper vector, 537
Russian wheat and Siberian oat mosaics compared, 539
seasonal occurrence and spread, 539
symptoms, 537
wound-tumor, filterability, 390
insect vectors, 9
pH and stability, 389
potato yellow-dwarf, stability compared, 390
properties of, 387
tomato spotted wilt, stability compared, 390
Vitamins, see also Nutrition, B_{12} absence in cedar-apple rust galls, 362
biotin effect on growth of *Chalara quercina*, 443
thiamine effect on growth of *Chalara quercina*, 443
Vitis spp., see also Grape, Melanconium leaf and stem fleck on, 347
Vivotoxins, as competitive inhibitors, 232
defined, criteria for, 229
mechanisms of action, vascular occlusion, 231
Volcani, Zafrira, 92

Wadsworth, D. F., 294, (294)
Waggoner, P. E., (195), (229), 281, 317, (319), (470), 488, (619), (663)
Waite, B. H., (700)

Walker, J. C., (116), 215, (415), (489), (490), (535), (596), 649
Wallin, J. R., 505
Walter, James M., (344)
Walters, H. J., (483)
Watermelon, anthracnose, breeding for resistance to, 254
cucumber Fusarium on, 481
Fusarium wilt, breeding for resistance to, 254
Watertor, Jean L., (475)
Watkins, G. M., (291)
Watson, R. D., 590
Weather, effect on production and survival of sporangia of *Phytophthora infestans*, 506
jelly end rot of potato effect on, 467
microclimate effect on white pine blister rust distribution, 487
Weathers, Lewis G., (482), 488, (550), (669)
Weaver, L. O., 407
Weber, George F., 488
Weed, Richard M., (480)
Weihing, John L., 488
Weinberger, J. H., (691)
Weintraub, M., (175), (324), 328
Weiss, Freeman, 407
Wheat, black stem rust, races of pathogen manner of infection of resistant varieties, 485
bunt, conditions affecting resistance, 473
control, 393
fungi, survival in mixture on host, 558
pathogenicity of physiologic races and inbred lines of pathogen, 398
cereal yellow-dwarf virus, aphid vectors, 309, **312**
epiphytotic, conditions favoring, 310
host range, 309
crown root infection and rootrot, 484
crown rot, Hessian fly relation to, 467
disease distribution in Oklahoma, 294
ergot associated with loose smut, 461
foot rot, Hessian fly relation to, 467
Helminthosporium spp., reactions to isolates of, 517
Kubanka, *Puccinia graminis tritici*, effect of light and temperature on reaction to races of, 481
powdery mildew, seedling varieties in greenhouse, resistance, 290
temperature effect on, in greenhouse, 290
rootrot, 484
control, biological, new method, 701
Russian mosaic virus and striate mosaic compared, 539
rust, breeding for resistance, "composite variety," 467
rust, leaf, control by and phytotoxicity of chemotherapeutants, 496
distribution in Oklahoma, 294
races, 294
rust, stem, breeding for resistance, 472
chemotherapy, 659
control by and phytotoxicity of chemotherapeutants, 496
races, pathogenicity on Kenya varieties, 474
seedling and adult changes in resistance, 468
spore shower, origin and extent, 471
smut, covered, control, 461
smut, loose, control, seed treatment, 315
distribution in Oklahoma, 294
ergot associated with, 461
races of pathogen on varieties, 682
races, spore germination characteristics, 482
resistance of Kawvale, to, 683
Wheat—*continued*
speckled leaf blotch, distribution in Oklahoma, 294
streak mosaic virus, *Aceria tulipae* vector, 485
effect of wind and blowing soil on infection, 484
striate mosaic virus, host range, 539
leafhopper vector, 537
Russan wheat and Siberian oat mosaics compared, 539
seasonal occurrence and spread, 539
symptoms, 537
varietal reactions to diseases in Oklahoma, 294
yellow-dwarf virus, 131
effect on yield, 135
yellow rust, races, inheritance of resistance, 471
Wheeler, H. E., 236, 489
Wiant, James Stewart, Biographical sketch, 592
Wilbur, Wesley D., (469)
Wildfire toxin, 232
Wildman, S. G., (587), (589)
Wiles, A. B., (215), 489
Wilhelm, Stephen, (589), 590, 593
Wilkinson, R. E., 489
Williams, Edwin B., (483)
Williams, Lansing E., 489
Williamson, C. E., 489
Willison, R. S., 175, 324, (328)
Wilson, J. D., 489
Wilson, R. A., (405)
Winstead, N. N., 489, 490
Wolf, Frederick T., 294
Wollman, E. S. H., 490
Wood decay in buildings, 20
Wood rot, citrus, concentric canker, 99
Fomes applanatus, associated with, on citrus, 99
Woody tissue, instruments for isolation of fungi, 403
Wooley, Paul H., (589), 590
Wound cork, Fusarium dry rot of potatoes, effect of, on penetration of, 95
Wound hormones, 352
Wound periderm and cuticle, gladiolus corms, effect of heat-curing on, 141

Xanthomonas pelargonii on geranium, 588
phaseoli var. *sojensis*, field and laboratory tests on persistence, 189
pruni, bacteriophage, factors affecting plaque formation, 27
terramycin control on peach, 405
Xiphinema americanum, see Nematodes, dagger

Yale, John W., Jr., (590), 590
Yarwood, C. E., 70, 490, 590, 675
Young, Donald A., 490
Young, H. C., Jr., (294), 294

Zaumeyer, W. J., (11), 38, 45, (333), 407, (480)
Zentmyer, George A., 490
Zinc dimethyl dithiocarbamate, see Fungicides, ziram
ethylenebisdithiocarbamate, see Fungicides, zineb
Zinnia, cucumber mosaic virus strains on, cross-protection, 336
Zinnia elegans, nasturtium mosaic virus on, 306
Zscheile, F. P., (473)
Zuckerman, Bert M., (61), (65), 287, 490
Zygophiala jamaicensis, carnation greasy spot, 585
Microthyriella rubi, relation to, 470